



Clinical Practice Guideline

Post-Operative Care in the Kidney Transplant Recipient

Final Version:	February 2017
Review Date:	February 2022

Authors:

Dr Richard J Baker

Consultant Nephrologist, Renal Unit, St. James's University Hospital, Leeds

Dr Patrick B Mark

Clinical Reader in Nephrology, Honorary Consultant, Glasgow Renal and Transplant Unit, Queen Elizabeth University Hospital, Glasgow

Dr Rajan K Patel

Consultant Nephrologist, Glasgow Renal and Transplant Unit, Queen Elizabeth University Hospital, Glasgow

Dr Kate K Stevens

Clinical Lecturer in Nephrology, Glasgow Renal and Transplant Unit, Queen Elizabeth University Hospital, Glasgow

Nicholas Palmer

British Kidney Patient Association

Endorsements



The National Institute for Health and Care Excellence (NICE) has accredited the process used by the Renal Association to produce its Clinical Practice Guidelines. Accreditation is valid for 5 years from January 2017. More information on accreditation can be viewed at www.nice.org.uk/accreditation



This guideline has also been endorsed by the British Transplantation Society.

Method used to arrive at a recommendation

The recommendations for the first draft of this guideline resulted from a collective decision reached by informal discussion by the authors and, whenever necessary, with input from the Chair of the Clinical Practice Guidelines Committee. If no agreement had been reached on the appropriate grading of a recommendation, a vote would have been held and the majority opinion carried. However this was not necessary for this guideline.

Conflicts of Interest Statement

All authors made declarations of interest in line with the policy in the Renal Association Clinical Practice Guidelines Development Manual. Further details can be obtained on request from the Renal Association.

Table of Contents

1. Introduction	4
2. Summary of Clinical Practice Guidelines for the Post-Operative Care of the Kidney Transplant Recipient	5
3. Summary of Audit Measures for the Post-operative Care of the Kidney Transplant Recipient	18
4. Rationale for Clinical Practice Guidelines for Post-Operative Care of the Kidney Transplant Recipient	21
5. Lay Summary	72
6. Acknowledgements	73
Appendix	73

1. Introduction

This document is intended for those engaged in the care of kidney transplant recipients (KTR) who are non-experts. With increasing efforts to deliver health care locally, many renal transplant recipients are followed up in centres remote from the main surgical transplant unit. At the same time, transplantation medicine has evolved into an increasingly complex and specialised field of nephrology. The following guidelines reflect this alteration in clinical practice and are intended for those healthcare professionals who look after renal transplant patients. They are also intended to be useful to both medical and surgical trainees, general practitioners, nurse specialists and other associated healthcare professionals involved in the care of renal transplant patients.

These guidelines cover the period after renal transplantation, specifically from initial hospital discharge until graft failure or patient death. The management of KTR can be divided into two phases:

- a. an early post-operative phase when prevention of acute rejection, optimization of graft function and prevention of opportunistic infection are paramount
- b. a later phase when the aims are to preserve good graft function, ensure adherence to medication, and prevent the long-term consequences of immunosuppression – malignancy, infection and premature cardiovascular disease.

The transition between these two phases occurs around 3-6 months after transplantation at the time when the progressive, protocolised reduction in immunosuppression following transplantation reaches long-term maintenance levels. Management of the early and late phase complications of transplantation requires monitoring at reducing frequency, awareness of complications, access to investigation, and strategies for prevention and treatment of complications (ranging from early acute rejection, to late cardiovascular disease). There are regional differences in demographics, risk and organisation of services. The priority is agreement of local strategies for post-transplant management.

These guidelines are designed to complement those previously published relating to pre-transplant care.

It should be noted that other comprehensive guidelines have been published and reference will be made to these¹. In keeping with other guidelines issued by the Renal Association, we have used the modified GRADE system. This grading system classifies expert recommendations as 'strong' (Grade 1) or 'weak' (Grade 2) based upon balance between the benefits and risks, burden and cost. The quality or level of evidence is designated as high (Grade A), moderate (Grade B), low (Grade C) or very low (D) depending on factors such as study design, directness of evidence and consistency of results. Grades of recommendation and quality of evidence may range from 1A to 2D^{2,3}.

These guidelines are based upon systematic literature searches conducted between November 2014 and February 2016. The main searches were performed in November-December 2014 and then rerun in February 2016. We searched Pubmed/MEDLINE, the Cochrane database of systematic reviews and hand searched reference lists and articles identified by the writing group members up till March 2016. We also reviewed all related guidelines from the National Institute for Clinical Excellence, NHS Blood and Transplant, the Advisory Committee on the Safety of Blood, Tissues and Organs (SaBTO), Kidney Disease Improving Global outcomes (KDIGO), the European Renal Association Best Practice Guidelines, Caring for Australians with Renal Impairment (CARI) and American Society of Transplant Surgeons. We cross-referenced with the previous iteration of these guidelines. The Pubmed search terms used were 'kidney transplant' AND rejection/ rejection/ immunosuppression/ cancer/ cardiovascular/ diabetes/ obesity /smoking/ hypertension/ pregnancy/ gout/ infection/ vaccination/ fertility/ pregnancy.

References

1. Kidney Disease: Improving Global Outcomes Transplant Work G. KDIGO clinical practice guideline for the care of kidney transplant recipients. American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons 2009;9 Suppl 3:S1-155.
2. Mactier R, Davies S, Dudley C, et al. Summary of the 5th edition of the Renal Association Clinical Practice Guidelines (2009-2012). Nephron Clin Pract 2011;118 Suppl 1:c27-70.
3. Uhlig K, Macleod A, Craig J, et al. Grading evidence and recommendations for clinical practice guidelines in nephrology. A position statement from Kidney Disease: Improving Global Outcomes (KDIGO). Kidney Int 2006;70:2058-65.

2. Summary of Clinical Practice Guidelines for the Post-Operative Care of the Kidney Transplant Recipient

1. Kidney Transplant Recipient (KTR): Organisation of Outpatient Follow-up (Guidelines 1.1–1.4)

Guideline 1.1 – KTR: Clinic infrastructure

We suggest that the following infrastructure should be in place for KTR follow up (2D):

- A consultant-level health care professional should be available for every transplant clinic
- KTRs should be reviewed in a dedicated outpatient area
- The results of blood tests (including drug levels if possible) should be available within 24 hours
- A formal mechanism should exist for results review by health care professionals within 24 hours of a clinic appointment
- There should be access to a multidisciplinary renal team including pharmacist, dietician, social worker and psychologist
- Patient care should be planned along principles set out in the National Service Framework and “Kidney Health Delivering Excellence”

Guideline 1.2 – KTR: Clinic frequency

We suggest that uncomplicated patients may be reviewed progressively less frequently (2C)

- 2-3 times weekly for the first month after transplantation
- 1-2 times weekly for months 2-3
- Every 2-4 weeks for months 4-6
- Every 4-6 weeks for months 6-12
- 3-6 monthly thereafter

Guideline 1.3 – KTR: Patient access

We suggest that all patients should have access to support services and results. (2C)

- All patients should have the option of on-line access to their results via the “Patient View” service
- All patients should have open access to the renal transplant outpatient service and have an established point of contact for enquiries
- Patient information should be available in both written and electronic formats

Guideline 1.4 – KTR: Chronic transplant care review

We suggest that a detailed review should be performed annually post-operatively (2C)

- A process should exist for patient review on an annual basis in a different format of clinic according to the “Care plan model”
- This should be a patient-centred clinic, facilitated by a health care professional
- It should address concerns in medical, social, psychological and sexual domains
- Access to a renal dietician, social worker, specialist renal pharmacist and/or psychologist should be readily available from this clinic
- This process should proceed in parallel with formal medical review

2. Kidney Transplant Recipient: Non-adherence (Guideline 2.1)

Guideline 2.1 – KTR: Recognising non-adherence

We suggest that it is important to prevent and detect non-adherence in kidney transplant recipients. (2C)

- Factors associated with non-adherence should be identified
- An established interventional pathway should be in place for those at high risk of or with proven non-adherence
- Pathways should be in place for paediatric KTRs in transition and for adolescent KTRs

3. Kidney Transplant Recipient: Immunosuppressive treatment (Guidelines 3.1–3.12)

Guideline 3.1 – KTR: immunosuppression regimen

We recommend that the patient and/or carer should be engaged in the decisions around selection of induction agent and maintenance immunosuppression (1D)

Guideline 3.2 – KTR: Induction immunosuppression

We recommend induction therapy should take into account the following:

- Immunosuppressive drugs should be started before or at the time of renal transplantation (1B)
- Induction therapy with biological agents should be administered to all KTRs. In patients at low immunological risk this will generally involve an interleukin-2 receptor antagonist (IL2-RA). Recipients at higher immunological risk may be considered for T-cell (lymphocyte) depleting antibodies (TDAs)(1B)
- Induction therapy with TDAs may also be useful for lower immunological risk patients with the intention of either steroid or calcineurin inhibitor (CNI) avoidance (1C)

Guideline 3.3 – KTR: Induction immunosuppression

We suggest that a CNI should be started at the time of transplantation and not delayed until the graft is functioning (2C)

Guideline 3.4 – KTR: Maintenance immunosuppression

We recommend that maintenance immunosuppression should normally consist of a calcineurin inhibitor (CNI) and an anti-proliferative agent, with or without corticosteroids in low and medium immunological risk KTRs (1B)

Guideline 3.4 – KTR: Maintenance immunosuppression

We suggest that low-medium dose tacrolimus (trough target 4-8 ng/mL) is recommended as the CNI of choice in patients also taking steroids who are low and medium immunological risk and are not at high risk of developing post transplant diabetes mellitus (PTDM) (2C)

Guideline 3.5 – KTR: Maintenance immunosuppression

We suggest that mycophenolic acid -based drugs should be the first-line anti-proliferative agent in preference to azathioprine, except in fertile KTRs who are unwilling to use reliable contraception (2B)

Guideline 3.5 – KTR : Maintenance immunosuppression

We suggest that slow release tacrolimus may be used as an option as second line agents for patients who suffer intolerable side effects related to peak dose toxicity (2C)

Guideline 3.6 – KTR : Maintenance immunosuppression

We suggest that KTRs who are unable to tolerate tacrolimus or who suffer serious adverse reactions related to its use be considered for the use of second line agents such as ciclosporin, sirolimus, everolimus, or belatacept (1B)

Guideline 3.7 – KTR : Maintenance immunosuppression

We suggest that MPA-based drugs should be the first-line antiproliferative agent, in preference to azathioprine, except in fertile KTRs who are unwilling to use reliable contraception (2B)

Guideline 3.8 – KTR: Maintenance immunosuppression

We suggest that mycophenolate mofetil (MMF) and enteric-coated mycophenolate sodium (Myfortic[®]) provide equivalent maintenance immunosuppression (2B)

Guideline 3.9 – KTR: Maintenance immunosuppression

We suggest that steroid avoidance or steroid withdrawal can be used during the first week after transplantation in low immunological risk kidney transplant recipients (2B)

Guideline 3.10 – KTR: Maintenance immunosuppression

We suggest aiming for minimum target levels for CNIs in uncomplicated renal transplantation after 3 months (2C)

Guideline 3.11 – KTR: Maintenance immunosuppression

We suggest that CNIs should not be withdrawn (2B)

Guideline 3.12 – KTR: Maintenance immunosuppression

We suggest that if steroids are not withdrawn within the first month, then they should be continued at low dose (prednisolone 5 mg per day or less) (2C)

Guideline 3.13 – KTR: Monitoring of immunosuppression

We suggest that long-term monitoring of immunosuppression levels is required as follows:

- Tacrolimus and ciclosporin levels should be monitored. The initial frequency should be three times a week. Levels should also be checked when any medication with possible interactions is prescribed, the dosage is changed, the formulation is changed, or when there is unexplained graft dysfunction (2C)
- Tacrolimus should be monitored by the trough (C_0) level, while ciclosporin can be monitored by either C_0 or 2-hour post dose (C_2) level (2C)
- Tacrolimus and ciclosporin levels should be available within 24 hours of taking blood samples in the first three months after transplantation (2D)
- The utility of monitoring mycophenolic acid (MPA) C_0 levels is uncertain (2D)
- Sirolimus should be monitored by the C_0 trough level (2C)

Guideline 3.14 – KTR: Prescribing and the use of generic agents

We suggest that generic immunosuppression compounds should not be used unless they have been shown to be bioequivalent and approved by the European Agency for the Evaluation of Medicinal Products (2D)

Guideline 3.15 – KTR: Prescribing and the use of generic agents

We suggest that KTRs should be made aware of the existence of generics and the importance of not switching between preparations without appropriate supervision (2D)

Guideline 3.16 – KTR: Prescribing and the use of generic agents

We suggest that drugs should be prescribed by brand name (whether branded or generic drugs are prescribed) (2D)

Guideline 3.17 – KTR: Prescribing and the use of generic agents

We suggest that KTRs should be closely monitored after switching between generic preparations until a new steady state is established (2D)

4. Kidney Transplant Recipient: Acute rejection (Guidelines 4.1–4.12)

Guideline 4.1 – KTR: Diagnosis of acute rejection

We recommend that a transplant renal biopsy should be carried out before treating an acute rejection episode unless this will substantially delay treatment or pose a significant risk to the patient (1C)

Guideline 4.2 – KTR: Diagnosis of acute rejection

We suggest that two cores of renal tissue should be obtained at transplant biopsy since this will increase the sensitivity of the investigation (2C)

Guideline 4.3 – KTR: Diagnosis of acute rejection

We suggest that a 16 gauge automated core biopsy needle is used where possible to provide the best compromise between diagnostic usefulness and patient tolerance of the procedure (1C)

Guideline 4.4 – KTR: Diagnosis of acute rejection

We recommend that a protocol transplant renal biopsy, defined as a biopsy performed in a stable graft without clinical evidence of acute rejection, be considered in the setting of persisting delayed graft function (1C)

Guideline 4.5 – KTR: Diagnosis of acute rejection

We recommend that routine C4d and SV40 staining should be performed upon transplant biopsies to address other causes of graft dysfunction (2C)

Guideline 4.6 – KTR: Diagnosis of acute rejection

We suggest that a serum sample be sent at the time of renal biopsy (for graft dysfunction) to look for human leucocyte antigen (HLA)-specific antibodies (2C)

Guideline 4.7 – KTR: Treatment of acute rejection

We suggest that borderline acute cellular rejection should be treated in the context of acute graft dysfunction (2D)

Guideline 4.8 – KTR: Treatment of acute rejection

We recommend that high dose intravenous corticosteroids should be the first line treatment for acute cellular rejection (1D)

Guideline 4.9 – KTR: Treatment of acute rejection

We suggest that maintenance steroids should be added or restarted in steroid-free patients undergoing acute rejection of any type (2D)

Guideline 4.10 – KTR: Treatment of acute rejection

We suggest that lymphocyte depleting agents may be considered for refractory acute cellular rejection or aggressive vascular cellular rejection (i.e. Banff category 4 Type II and III) (2C)

Guideline 4.11.1 – KTR: Treatment of acute rejection

We suggest that antibody mediated rejection (AMR) should be treated with one or more of the following modalities: steroids; plasma exchange; intravenous immunoglobulin; anti-CD20 antibody, lymphocyte-depleting antibody or bortezomib (2C)

Guideline 4.11.2 -KTR: Treatment of acute rejection

We recommend that the British Transplant Society (BTS) guidelines on antibody incompatible transplantation for management of rejection in the context of antibody incompatible transplantation (1A-D)

Guideline 4.12 – KTR: Treatment of acute rejection

We suggest that - after an episode of rejection (unless associated with low CNI levels) - azathioprine should be switched to MPA-based immunosuppression, MPA should be started, or the existing dose of MPA maximised (2D)

5. Kidney Transplant Recipient: Chronic Allograft Injury (Guidelines 5.1–5.7)

Guideline 5.1 – KTR: Diagnosis of Chronic Allograft Injury

We recommend that early identification of graft injury is desirable to maximise the potential for intervention. A proactive and systematic approach should be employed to manage graft dysfunction (1C)

Guideline 5.2 – KTR: Detection of Chronic Allograft Injury

We suggest that renal function should be monitored at each clinic visit by assessment of serum creatinine and qualitative evaluation of urine protein excretion by dipstick, supplemented by spot protein:creatinine ratio (PCR) or albumin:creatinine ratio (ACR) if positive (2C)

Guideline 5.3 – KTR: Diagnosis of Chronic Allograft Injury

We suggest that renal biopsy is the optimal investigation for parenchymal causes of graft dysfunction where the cause is uncertain (2C)

Guideline 5.4 – KTR: Diagnosis of Chronic Allograft Injury

We suggest that renal biopsies in patients with chronically deteriorating function should be stained for C4d and SV40 (2C)

Guideline 5.5 – KTR: Diagnosis of Chronic Allograft Injury

We suggest that a serum sample should be sent at the time of renal biopsy (for graft dysfunction) to look for HLA-specific antibodies (2C)

Guideline 5.6 – KTR: Treatment of chronic allograft injury

We suggest that chronic allograft injury should be treated:

- By withdrawal of calcineurin inhibitors if there is histological evidence of CNI toxicity or non-specific interstitial fibrosis and tubular atrophy (2C)
- By intensification of immunosuppression if there is evidence of ongoing immune injury (cellular rejection and/or humoral rejection) (2C)
- In a similar fashion to other patients with chronic kidney disease (CKD), following similar preventative strategies and with timely referral to low clearance services (2D)

Guideline 5.7 – KTR: Renal biopsy in chronic allograft injury

We suggest that a renal transplant biopsy is indicated:

- If there is a persistent unexplained elevation of creatinine or failure to return to baseline after an episode of biopsy proven acute rejection (BPAR) (1C)
- Every 7-10 days during delayed graft function (DGF) (2C)
- If expected renal function is not achieved within 4-8 weeks (2D)
- If sustained new onset proteinuria develops (PCR >50 mg/mmol or ACR >35 mg/mmol) (2C)

6. Kidney Transplant Recipient: Cardiovascular Disease and Lifestyle (Guidelines 6.1–6.6)

Guideline 6.1 – KTR: Hypertension

We suggest that the management of hypertension take into account that:

- Blood pressure should be recorded at each clinic visit (1C)
- Clinic blood pressure should be <140/90 mmHg in clinic (130/80 mmHg if PCR >50 or ACR >35) (2C)
- Home blood pressure recordings and 24-hour ambulatory recordings may be helpful in some instances but lower BP targets should then be set (home and or ambulatory daytime measures <135/80mmHg) (2D)
- There is no evidence that any antihypertensive agent is better than any other and effort should be focused on achieving absolute blood pressure control rather than the use of individual agents (2D)
- Inhibitors of the renin-angiotensin system may be more effective in the minimisation of proteinuria but should be used with caution in the first 3 months post-transplant.(2C)
- Resistant hypertension may be due to transplant renal artery stenosis and should be investigated according to local practice (2D)

Guideline 6.2 – KTR: Dyslipidaemia

We suggest that the management of dyslipidaemia take into account that:

- Fasting lipid levels should be measured on an annual basis in renal transplant recipients (2C)
- Treatment targets should be the same as in the general population (2C)
- KTRs at increased primary or secondary cardiovascular risk receive statin therapy to reduce the risk of coronary artery disease (2C)
- The choice and dose of statin should take into account concurrent immunosuppression. High dose simvastatin (≥40mg daily) should be avoided in conjunction with ciclosporin and/calcium channel antagonists (2D)

Guideline 6.3 – KTR: Diabetes mellitus

We suggest that the detection and treatment of diabetes should consider:

- Screening for the development of post-transplant diabetes mellitus (PTDM) by dipstick urinalysis and measurement of blood sugar level at each clinic visit (2C)
- Post-transplant immunosuppression should take into account risk factors for the development of diabetes (2C)
- The diagnosis of PTDM is made based on WHO criteria for the diagnosis of diabetes mellitus based on fasting or random blood, serum glycated haemoglobin (HBA1c) or oral glucose tolerance testing (1C)
- A diagnosis of PTDM is made once patients are established on stable maintenance immunosuppression (2D)
- Post-transplant diabetes should be managed in collaboration with specialists in diabetic medicine (2D)
- All units should have a protocol for the management of post-transplant diabetes (2C)
- KTR with diabetes (either prior to transplantation or PTDM) should undergo screening for diabetic complications (retinal screening, foot care, neuropathy) in line with guidelines for non KTR patients with diabetes (2D)

Guideline 6.4 – KTR: Ischaemic heart disease

We suggest that KTRs receive standard treatment for ischaemic heart disease, including thrombolysis, revascularisation, and secondary prevention (2C)

Guideline 6.5 – KTR: Smoking cessation

We recommend that smoking should be strongly discouraged in transplant recipients (see guideline 6.4) (1A)

Guideline 6.6 – KTR: Lifestyle measures

We suggest that advice on healthy lifestyle forms a routine part of post-transplant care:

- Maintenance of a healthy diet should be encouraged (2C)
- An ideal weight should be targeted (body mass index (BMI) ≤ 25 kg/m²) (2C)
- Weight management services should be available (2C)
- We suggest that KTRs participate in physical activity at a level similar to that recommended to age and co-morbidity matched counterparts from the general population (2D)
- Alcohol consumption should be within national guidelines (2D)
- Recreational drug use should be avoided (2D)
- The use of over-the-counter medications (without discussion with clinical staff) and non-proprietary medications (e.g. herbal medicines) should be discouraged (2D)

7. Kidney Transplant Recipient: Neoplasia (Guidelines 7.1–7.4)

Guideline 7.1 – KTR: Screening for cancer

We suggest that the organisation of screening for neoplasia in KTRs take into account:

- Screening should be similar to the general population for cervical, breast, colon and prostate cancer (2C)
- Screening is not recommended for renal cell carcinoma (2C)
- Patient education pre and post transplantation (1C)
- Patients should be aware of malignancy risk and encouraged to report symptoms which may represent *de novo* malignancy (e.g. breast or testicular lumps) (2D)
- Skin surveillance by a healthcare professional at least biannually up to 5 years post-transplant and annually from 5 years post-transplant (2C)
- Patients with cirrhosis should undergo an annual hepatic ultrasound and determination of serum alpha foeto-protein (2C)

Guideline 7.2 – KTR: Non-melanoma skin cancer (NMSC)

We recommend that KTRs should be educated about the adverse effects of sun exposure (1C)

Guideline 7.3 – KTR: Non-melanoma skin cancer

We suggest that KTRs that an individualised assessment of hazard should be made according to risk factors (2C)

Guideline 7.4 – KTR: Non-melanoma skin cancer

We recommend that patients should be encouraged to cover their skin in direct sunlight and to use total sunblock (Sun Protection Factor ≥ 50) (1D)

Guideline 7.5 – KTR: Non-melanoma skin cancer

We suggest that self-examination should be encouraged with guidance provided. This should be supplemented by at least biannual review by a trained healthcare professional up to 5 years post-transplant and annual review from 5 years (2C)

Guideline 7.6 – KTR: Non-melanoma skin cancer

We suggest that the prescription of acitretin as chemoprophylaxis be considered in those with ≥ 2 previous NMSC if there are no contraindications (2B)

Guideline 7.7 – KTR: Immunosuppression in cancers

We suggest that immunosuppression should be reduced if neoplasia develops (2C)

Guideline 7.8 – KTR: Immunosuppression in cancers

We suggest that mammalian target of rapamycin inhibitors (m-TORi) are considered as alternative immunosuppressive agents in KTRs who develop *de novo* malignancy (2C)

Guideline 7.9 – KTR: Immunosuppression in Kaposi sarcoma

We suggest that m-TORs have specific anti-tumour effects in Kaposi sarcoma (2C)

8. Kidney Transplant Recipient: Infection Complications (Guidelines 8.1–8.9)

Guideline 8.1 – KTR: Vaccination

Guideline 8.1.1 – KTR: Vaccination

We recommend that KTRs:

- Should be vaccinated with inactivated viruses as per the normal population (1D)
- Should receive annual influenza vaccination unless contraindicated (1C)

Guideline 8.1.2 – KTR: Vaccination

We suggest that KTRs:

- Should have hepatitis B surface antibody (HBsAb) levels rechecked annually and be revaccinated if antibody titres fall below 10 mIU/mL (2D)
- Should not receive live attenuated vaccines (2C)
- Should receive pneumococcal vaccine and a booster every five years (2D)

Guideline 8.2 – KTR: Cytomegalovirus (CMV) disease

Guideline 8.2.1 – KTR: Prophylaxis and treatment of CMV disease

We recommend:

- Prophylaxis should be continued for 3-6 months, until immunosuppression has been reduced to long-term maintenance level (1B)
- Treatment should be administered for 6 weeks after treatment with a TDA (1C)

Guideline 8.2.2 – KTR: prophylaxis and treatment of CMV disease

We suggest:

- All transplant units should be able to measure CMV serological status and quantify viral load (2D)
- Donor and recipient CMV status should be recorded at the time of transplantation (2D)
- Each unit should have a written protocolised CMV strategy based on prophylaxis or pre-emptive therapy (2D)
- For the treatment of mild and moderate CMV disease, oral valganciclovir and intravenous ganciclovir are of equivalent efficacy (2C)
- The first line treatment of life-threatening CMV disease is intravenous ganciclovir (2D)
- Treatment duration should be determined by monitoring viral load (2C)

Guideline 8.3 – KTR: Epstein Barr Virus (EBV) infection

Guideline 8.3.1 – KTR: EBV infection

We recommend that immunosuppression should be reduced or stopped following the development of post transplant lymphoproliferative disease (PTLD) (1C)

Guideline 8.3.2 – KTR: EBV infection

We suggest:

- Both donor and recipient should have their EBV serology recorded at the time of transplantation (2D)
- All high risk (D⁺/R⁻) patients (including adults) should have EBV viral load measured immediately after transplantation, monthly for six months, and three monthly to one year (2C)
- EBV viral load should be monitored after the treatment of rejection (2C)
- Total immunosuppression should be reduced when EBV titres rise significantly (2C)

Guideline 8.4 – KTR: Varicella Zoster Virus (VZV) infection

Guideline 8.4.1 – KTR: VZV infection

We recommend:

- Primary infection (chickenpox) should be treated with intravenous aciclovir or oral valaciclovir until the lesions scab over (1C)
- Uncomplicated shingles should be treated with oral acyclovir or valaciclovir until the lesions scab over (1D)
- Disseminated (>2 dermatomes), ocular or invasive shingles should be treated with intravenous aciclovir until the lesions scab over, together with a reduction in immunosuppression (1B)

- Varicella-susceptible KTRs (i.e. VZV IgG negative) with primary exposure to VZV should receive intravenous immunoglobulin, ideally within 96 hours, but up to a maximum of 10 days following exposure. If unavailable or after 10 days, oral aciclovir should be prescribed for seven days (1D)

Guideline 8.4.2 – KTR: VZV infection

We suggest:

- Patients on the waiting list who are VZV IgG negative should be vaccinated prior to transplantation (2D)
- Immunosuppression should be reduced during primary infection (2D)

Guideline 8.5 – KTR: Herpes Simplex Virus (HSV) infection

Guideline 8.5.1 – KTR: HSV infection

We recommend:

- Superficial HSV infection should be treated with appropriate oral agents until the lesions have resolved (1D)
- Systemic HSV infections should be treated with intravenous aciclovir and a reduction in immunosuppression until a response occurs and oral medication should be continued for at least 14 days (1C)

Guideline 8.5.2 – KTR: HSV infection

We suggest that KTRs suffering frequent recurrent HSV infection should consider oral prophylaxis (2D)

Guideline 8.6 – KTR: BK virus (BKV) nephropathy

Guideline 8.6.1 – KTR: BK nephropathy

We recommend that confirmed BK nephropathy should be treated by reduction in immunosuppression (1D)

Guideline 8.6.2 – KTR: BK nephropathy

We suggest:

- Screening should also be carried out when renal function deteriorates in an unexplained fashion (2D)
- KTRs should be screened for BKV viral load or by performing urine microscopy for decoy cells or by polymerase chain reaction (PCR) on urine or serum (2C)
- Suspected BK nephropathy should be confirmed by renal biopsy, which should be stained for SV40. Two cores containing medullary tissue should ideally be examined (2D)
- Immunosuppression should be reduced when the serum BKV load exceeds 10^4 copies/ml (2C)
- There is no established specific treatment for BK nephropathy (2D)
- Re-transplantation can safely be considered in patients who have BK nephropathy diagnosed in an earlier graft (2C)

Guideline 8.7 – KTR: Pneumocystis jirovecii infection - treatment and prophylaxis

We suggest:

- All patients with confirmation (microscopy or PCR) of Pneumocystis jirovecii in respiratory secretions should be treated for 14 to 21 days with co-trimoxazole orally or intravenously (15-20mg/kg in three or four divided doses) (2B)

- Patients with contraindications to treatment with co-trimoxazole should receive pentamidine (4mg/kg/day intravenously) (2B)
- Adjunctive glucocorticoid therapy may be considered in patients with severe disease (2D)
- All patients should receive 3-6 months of treatment with co-trimoxazole 480mg daily for *Pneumocystis jirovecii* prophylaxis following renal transplantation (1B)

Guideline 8.8 – KTR: Post-transplant infection prophylaxis

We suggest:

- All patients should receive 3-6 months of treatment with co-trimoxazole 480 mg daily (1B)
- Oral antifungal prophylaxis should be administered for one month after transplantation (2C)
- In selected patients, prophylaxis against mycobacterium tuberculosis with daily isoniazid (supplemented with pyridoxine) should be instituted for six months after transplantation (2C)

Guideline 8.9 – KTR: Hepatitis E Virus (HEV)

We recommend that Hepatitis E Virus (HEV)-screened blood components should be given to all KTR (1C)

9. Kidney Transplant Recipient: Bone and Joint Disease (Guidelines 9.1–9.4)

Guideline 9.1 – KTR: Osteoporosis

We suggest:

- KTRs suffering from osteoporosis or at high potential risk should be considered for steroid-avoiding immunosuppression (2D)
- KTRs on long-term steroids or at high risk for osteoporosis should undergo DEXA scanning if eGFR >30 mL/min/1.73m² (2D)
- Treatment should be according to the Royal College of Physicians (RCP) guidelines for steroid-induced osteoporosis (2D)

Guideline 9.2 – KTR: Tertiary hyperparathyroidism

We suggest:

- Severe hyperparathyroidism should be treated prior to transplantation (2D)
- Cinacalcet can be used in KTR (2C)
- Treatment should be the same as for other patients with CKD (2D)

Guideline 9.3 – KTR: Gout

Guideline 9.3.1 – KTR: Treatment of gout

- We recommend that neither allopurinol nor febuxostat should be administered with azathioprine (1C)

Guideline 9.3.2 – KTR: Treatment of gout

We suggest:

- Hyperuricaemia should be treated when associated with gout, tophi or uric acid stones (2D)
- Non-steroidal anti-inflammatory drugs (NSAIDs) should be avoided in KTRs (2D)
- Acute gout may be treated with brief a course of oral prednisolone. (2D)
- Colchicine is an effective treatment for gout in KTR (2D)

Guideline 9.4 – KTR: Calcineurin inhibitor bone pain

We suggest:

- Reduction or withdrawal of CNIs should be considered in KTRs with intractable bone pain (2D)
- Dihydropyridine calcium antagonists also may be beneficial (2D)

10. Kidney Transplant Recipient (KTR): Haematological Complications (Guidelines 10.1–10.3)

Guideline 10.1 – KTR: Anaemia

We suggest that chronic anaemia should be managed in the same way as other patients with CKD (2D)

Guideline 10.2 – KTR: Polycythaemia

We recommend that initial treatment should be with angiotensin converting enzyme inhibitors (ACEIs) or angiotensin receptor blockers (ARBs) (1C)

Guideline 10.3 – KTR: Polycythaemia

We suggest:

- Haemoglobin levels should be monitored at every clinic visit (2D)
- Treatment should be initiated if the haematocrit or packed cell volume exceeds 52% in men and 49% in women (2D)
- Venesection may be used in refractory cases.(2D)

11. Kidney Transplant Recipient (KTR): Reproductive Issues (Guidelines 11.1–11.5)

Guideline 11.1 – KTR: Conception and contraception (female)

We recommend that MPA-containing immunosuppressant drugs should be stopped prior to conception and replaced appropriately (1A)

Guideline 11.2 – KTR: Conception and contraception (female)

We suggest:

- KTRs should wait for one year after transplant and have stable function before attempting conception (2C)
- Counselling regarding fertility and reproduction should be offered to female KTRs and their partners either prior to transplantation or soon afterwards (2D)
- m-TORi should be stopped prior to conception and replaced as appropriate (2D)
- Pregnancy should be jointly managed with an Obstetrics department with experience of care of KTR (2D)

- KTRs receive aspirin 75 mg daily to reduce the risk of pre-eclampsia from 12 weeks gestation until birth of the baby unless there are contraindications (2C)
- The risks and benefits of breastfeeding should be discussed (2D)
- Contraception advice should be similar to the general population (2D)

Guideline 11.3 – KTR: Conception and contraception (male)

We recommend:

- Male KTRs are advised that MPA containing compounds have theoretical teratogenic potential in men taking these agents (1D)
- KTRs should be advised that m-TORi reduce the male sperm count and counselled accordingly. (1C)

Guideline 11.4 – KTR: Conception and contraception (male)

We suggest:

- All immunosuppressive drugs other than m-TORi can be used in male KTRs. Advice for MPA is as Guideline 11.3 (2D)
- The decision to continue MPA containing compounds in a male KTR wishing to conceive should balance the risk of theoretical teratogenicity against the risk of rejection on changing from MPA to azathioprine (2D)
- Men on m-TORi who wish to conceive should discontinue these agents prior to conception and replace them as appropriate (2D)
- Men who wish to maintain fertility should avoid m-TORi or bank sperm prior to starting these drugs (2D)

Guideline 11.5 – KTR: Sexual dysfunction

We suggest:

- Specific enquiry should be made regarding sexual dysfunction, preferably at an annual review clinic (2D)
- Care pathways for dealing with sexual dysfunction should be established (2D)
- Close liaison with local andrology service is recommended (2D)
- Sildenafil is safe and effective in male KTR not taking nitrates (2D)

3. Summary of Audit Measures for the Post-operative Care of the Kidney Transplant Recipient

- Proportion of blood results available for review, and reviewed, within 24 hours
- Proportion of units with a written follow-up schedule available to all staff and patients
- Percentage of patients accessing their results through Renal Patient View
- Percentage of total patients assessed in an annual review clinic
- Recording “Did Not Attend” (DNA) rates for all patients
- Recording sub-therapeutic drug levels. Measuring within-patient variability of CNI levels
- Percentage of total patients receiving induction with ILRAs and TDAs

- Percentage of de novo KTRs receiving tacrolimus
- Percentage of de novo KTRs receiving MPA based immunosuppression
- Percentage of de novo KTRs receiving corticosteroid maintenance therapy
- Use of generic agents
- Severity of biopsy proven acute rejection (BPAR) recorded by Banff criteria
- Percentage of KTRs with BPAR in first 3 months and first 12 months
- Percentage of KTRs requiring TDAs to treat rejection in first year
- Complication rates after renal transplant biopsy
- The percentage of KTRs with BPAR in first 3 months and the first 12 months
- The percentage of KTRs with a donor specific HLA antibody at the time of biopsy
- The percentage of KTRs with positive C4d staining on biopsy
- Proportion of patients receiving a target blood pressure of 140/90 mmHg or 130/80 mmHg in the presence of proteinuria (PCR >100 mg/mmol or ACR >70 mg/mmol)
- Proportion of patients with proteinuria assessed by dipstix and, if present, quantified at each clinic visit
- Proportion of renal transplant recipients with an annual fasting lipid profile
- Proportion of RTR taking statins (including the type of statin) for primary and secondary prevention of premature cardiovascular disease
- Proportion of patients on other lipid lowering agents
- Proportion of patients achieving dyslipidaemia targets
- Incidence of post-transplant diabetes mellitus (PTDM) at three months and at annual intervals thereafter
- Proportion of patients who require insulin, and in whom remedial action is undertaken – minimisation of steroids and switching of CNIs
- The proportion of patients with PTDM enrolled in screening for extra-renal complications of PTDM
- Proportion of patients with ischaemic heart disease
- Proportion of patients suffering myocardial infarction
- Proportion of patients undergoing primary revascularisation

- Proportion of patients receiving secondary prevention with a statin, anti-platelet agents and renin angiotensin system (RAS) blockers
- Proportion of KTRs who smoke
- Proportion of cigarette smoking KTRs who have been given formal advice or offered help with cessation
- Proportion of patients who are obese (BMI>30 kg/m²)
- Proportion of patients having screening procedures for neoplasia at the annual review clinic
- Incidence of CMV disease
- Rate of EBV infection and PTLD
- Completeness of records for EBV donor and recipient serology
- Rates of primary VZV and shingles infection
- Completeness of records for VZV recipient serology
- Rates and outcomes of HSV infections
- Rates of BK viral infection in screening tests
- Rates and outcomes of BK nephropathy
- Frequency of bisphosphonate use
- Incidence of fractures
- Incidence of hyperparathyroidism
- Incidence of parathyroidectomy
- Use of cinacalcet
- Frequency of hyperuricaemia and gout
- Prevalence of anaemia
- Prevalence of polycythaemia
- Pregnancy rates and outcomes
- Prevalence of sexual dysfunction

4. Rationale for Clinical Practice Guidelines for Post-Operative Care of the Kidney Transplant Recipient

1. Kidney Transplant Recipient (KTR): Organisation of Outpatient Follow-up (Guidelines 1.1 – 1.4)

Guideline 1.1 – KTR: Clinic infrastructure

We suggest that the following infrastructure should be in place for KTR follow up (2D)

- A consultant-level health care professional should be available for every transplant clinic
- KTRs should be reviewed in a dedicated outpatient area
- The results of blood tests (including drug levels if possible) should be available within 24 hours.
- A formal mechanism should exist for results review by health care professionals within 24 hours of a clinic appointment
- There should be access to a multidisciplinary renal team including pharmacist, dietician, social worker and psychologist
- Patient care should be planned along principles set out in the National Service Framework and “Kidney Health Delivering Excellence”

Audit Measure

The proportion of blood results available for review, and reviewed, within 24 hours

Rationale

All KTRs should have ready access to a senior clinical opinion and a senior clinician should be available at renal transplant clinics. In some centres this may be a consultant-level nurse, in others a medical or surgical consultant. The exact type of healthcare professional is not important but KTRs and junior staff should have access to an individual with appropriate knowledge and experience. This will also benefit the training of junior medical staff. A dedicated outpatient area is beneficial as it provides a familiar environment and staff experienced in the management of patients on renal replacement therapy.

Prompt availability and formal review of test results is desirable since most complications can be resolved more easily if recognised at an early stage, particularly in the first few weeks after renal transplantation. It is recommended that patient care is carried out according to the principles laid out in the Department of Health (DoH) leaflet, “Achieving Excellence in Kidney Care”⁴ and the report by Kidney Research UK “Kidney Health Delivering Excellence”⁵.

References

2. Health Do. Achieving Excellence in Kidney Care: Delivering the National Service Framework for Renal Services. 2009.
3. UK KR. Kidney Health Delivering Excellence. 2013.

Guideline 1.2 – KTR: Clinic frequency

We suggest that uncomplicated patients, in general, may be reviewed progressively less frequently in clinic (2C)

- 2-3 times weekly for the first month after transplantation
- 1-2 times weekly for months 2-3
- Every 2-4 weeks for months 4-6
- Every 4-6 weeks for months 6-12
- 3-6 monthly thereafter

Audit Measure

Proportion of units with a written follow-up schedule available to all staff and patients

Rationale

Freedom from regular hospital attendance is an important benefit of renal transplantation, balanced against the risks and prevention of complications. These risks (especially of surgical complications) are highest in the immediate postoperative period and during the first few weeks following hospital discharge, when the burden of immunosuppression is greatest. For typical patients monitoring should therefore be most frequent during this period and then diminish with time. The use of virtual renal clinics should be explored as a complementary form of KTR review as it might be more convenient for some patients.

Guideline 1.3 – KTR: Patient access

We suggest that all patients should have ready access to support services and results (2C)

- All patients should have on-line access to their results via the “Renal Patient View” service if they wish
- All patients should have open access to the renal transplant outpatient service and have an established point of contact for enquiries
- Patient information should be available in both written and electronic formats

Audit Measure

Percentage of patients accessing their results through Renal Patient View

Rationale

Patients should be encouraged to take an active role in their own care according to principles embodied in the National Service Framework⁴. Interest in their own blood results should be welcomed and KTRs should be encouraged to use Patient View (<https://www.patientview.org/#/>; formerly known as Renal Patient View). Patient education is a crucial element in the success of renal transplantation and easy access to information should be provided for all patients in different formats (e.g. paper-based and electronic).

References

4. Health Do. Achieving Excellence in Kidney Care: Delivering the National Service Framework for Renal Services. 2009.

Guideline 1.4 – KTR: Chronic transplant care review

We suggest that a detailed review should be performed annually post-operatively (2C)

- A process should exist for patient review on an annual basis in a different format of clinic according to the “Care plan model”
- This should be a patient-centred clinic, facilitated by a health care professional
- It should address concerns in medical, social, psychological and sexual domains
- Open access to a renal dietician, social worker, specialist renal pharmacist and/or psychologist should be readily available from this clinic
- This process should proceed in parallel with formal medical review

Audit Measure

Percentage of total patients assessed in an annual review clinic

Rationale

Since KTRs experience considerable late morbidity which is unlikely to be managed properly in a traditional clinical setting (e.g. skin lesions, sexual dysfunction and psychological morbidity) it seems sensible to facilitate periodic follow up in a different and more holistic environment^{6,7}.

References

4. Reece SM, Harden PN, Smith AG, Ramsay HM. A model for nurse-led skin cancer surveillance following renal transplantation. *Nephrology nursing journal : journal of the American Nephrology Nurses' Association* 2002;29:257-9, 67.
5. Short CD, Russell S, Valentine A. Clinical audit and long-term evaluation of renal transplant recipients. *Transplantation* 2001;72:S94-8.

2. Kidney Transplant Recipient (KTR): Non-adherence (Guideline 2.1)

Guideline 2.1 – KTR: Recognising non-adherence

We suggest that it is important to prevent and detect non-adherence in kidney transplant recipients (2C)

- Factors associated with non-adherence should be identified
- An established interventional pathway should be in place for those at high risk of or with proven non-adherence
- Pathways should be in place for paediatric KTRs in transition and for adolescent KTRs

Audit Measures

- Recording “Did Not Attend” (DNA) rates for all patients
- Recording sub-therapeutic drug levels
- Measuring within-patient variability of CNI levels

Rationale

Non-adherence with immunosuppressive medication is an important factor in graft loss and up to a third of patients report regularly missing tablets⁸⁻¹¹. One Dutch study described self-reported non-adherence rates of 17% at six weeks after transplantation rising to 27% by six months¹². Clinical parameters associated with non-adherence are well recognised and should be used to assessing risk e.g. erratic or low immunosuppression levels, clinic non-attendance, psychiatric illness, low belief in the need for medication, adolescence and early adulthood⁸⁻¹⁰. The patient and/or their carer should be fully engaged in identifying reasons for and strategies to address non-adherence. High within-patient variability of CNI levels has also been shown to be associated with poor graft outcomes and can readily be monitored in the clinic¹³⁻¹⁵. It remains to be proven whether there is a prospective intervention in such patients that can improve outcomes.

References

8. O'Grady JG, Asderakis A, Bradley R, et al. Multidisciplinary insights into optimizing adherence after solid organ transplantation. *Transplantation* 2010;89:627-32.
9. Prendergast MB, Gaston RS. Optimizing medication adherence: an ongoing opportunity to improve outcomes after kidney transplantation. *Clinical journal of the American Society of Nephrology : CJASN* 2010;5:1305-11.
10. Tong A, Howell M, Wong G, Webster AC, Howard K, Craig JC. The perspectives of kidney transplant recipients on medicine taking: a systematic review of qualitative studies. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2011;26:344-54.
11. Butler JA, Peveler RC, Roderick P, Smith PW, Horne R, Mason JC. Modifiable risk factors for non-adherence to immunosuppressants in renal transplant recipients: a cross-sectional study. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2004;19:3144-9.
12. Massey EK, Tielen M, Laging M, et al. The role of goal cognitions, illness perceptions and treatment beliefs in self-reported adherence after kidney transplantation: a cohort study. *Journal of psychosomatic research* 2013;75:229-34.
13. Kahan BD, Welsh M, Urbauer DL, et al. Low intraindividual variability of cyclosporin A exposure reduces chronic rejection incidence and health care costs. *Journal of the American Society of Nephrology : JASN* 2000;11:1122-31.
14. Sapir-Pichhadze R, Wang Y, Famure O, Li Y, Kim SJ. Time-dependent variability in tacrolimus trough blood levels is a risk factor for late kidney transplant failure. *Kidney Int* 2014;85:1404-11.
15. van Gelder T. Within-patient variability in immunosuppressive drug exposure as a predictor for poor outcome after transplantation. *Kidney Int* 2014;85:1267-8.

3. Kidney Transplant Recipient (KTR): Immunosuppressive treatment (Guidelines 3.1 – 3.17)

General concepts

The starting point for renal transplantation is comparison with other forms of renal replacement therapy (RRT). Renal transplantation provides a better quality of life, an increased sense of well-being and a longer life span when compared to other forms of RRT. Therefore minor differences in clinical outcome between different immunosuppressive regimes should be placed in context with the much greater difference in outcome between transplantation and other forms of RRT, for those fit enough to be wait listed (approximately 30% of those with end stage renal disease).

Almost all renal transplants are allogeneic (i.e. not from identical twins) and will provoke a powerful immunological rejection response in the recipient. Rejection will destroy renal tissue and so the primary aim of immunosuppression is to avoid rejection. In general, more potent immunosuppressive regimes will reduce the risk of all forms of rejection but at the expense of increased side effects. Side effects comprise generic immunosuppressive side effects (e.g. increased risk of infection and malignancy) or those specific to the particular drug used (e.g. gingival hypertrophy with ciclosporin).

Immunosuppressive management may be divided into three phases – induction, early (<3-6 months post-transplant), and late (>3-6 months). More intensive immunosuppression is required in the early post-operative period to prevent acute rejection episodes, while long-term immunosuppression should balance the risk of rejection against the adverse effects of immunosuppressive therapy. Effective immunosuppression is best achieved by combination therapy that minimises the side effects of individual agents. Overall, the aim of immunosuppression is to maximise patient and graft survival following transplantation and to maximise the quality of life and economic benefits of transplantation.

When planning immunosuppressive treatment, it is essential to consider the risks to the recipient. The risks of immunosuppressive therapy are largely predictable and should be balanced against the risk of harm to the individual patient from under-immunosuppression and resulting rejection, and the benefits of a well-functioning transplant. The assessment of risk is imprecise but Table 1 illustrates some broad principles. Many transplant units employ such risk stratification but there is very little evidence to support such an approach since most studies have excluded high-risk patients.

Risk Type	Low	Medium	High	Possible Strategy
Immunological	0-DR mismatch First graft Unsensitised Recipient >60	1-DR mismatch Afro-Caribbean recipient Historical DSAs NDSAs DGF Older donor ¹⁶	2-DR mismatch Previous early immunological graft loss DSAs ABO-incompatible Sensitised (high CRF/PRA) Preoperative anti-ATIIR Abs ¹⁷	Increase total immunosuppressive load
Metabolic	Low BMI Age <40 Normal pre-Tx GTT	Positive family history ADPKD	Impaired GT BMI >35 HCV positive Age >60 Previous CVD Race	Avoid/minimise steroids and tacrolimus
Neoplastic	Age <40	Pre-malignant lesion	Previous cancer Hereditary syndrome e.g. VHL	Consider low immunosuppression load or sirolimus
Ischaemia-reperfusion injury	Living donor Deceased donor <40	CIT >12 hours Donor aged 50-60	DCD CIT > 24 hours Extended criteria donor	Reduce CNI exposure
Non-adherence			Poor RRT compliance Age <20 Transition from paediatric to adult	Education Simple drug regime alemtuzumab or belatacept

Table 1. Risk stratification for selection of immunosuppression in kidney transplantation

For the purpose of these guidelines, immunosuppression has been broadly divided into induction and maintenance phases; the maintenance phase can be further divided into early and late. While the distinction between these periods is largely arbitrary, here the induction period is considered as the peri-transplant period, the early maintenance period is the 3-6 month period after transplantation when immunosuppression is tapered, and the late maintenance phase is the period beyond 3-6 months when immunosuppression has been tapered to long-term levels. It is recognised that the renal allograft is more immunogenic during the early post-transplant period and that more potent immunosuppression is therefore required to prevent rejection. In the later maintenance phase the allograft becomes less immunogenic and more consideration can be given to the minimisation of side effects from immunosuppression.

Immunosuppressive strategies may be pre-emptive or reactive. For example, steroid avoidance is pursued by some units with the objective of avoiding steroid-related side effects. It also permits the widespread usage of tacrolimus with a reduced risk of post-transplant diabetes mellitus (PTDM). Other units adopt a strategy of dual or lower dose triple immunosuppressive drug therapy.

When considering the published evidence, it is essential to look at long-term outcome data. However, long-term data from adequately powered clinical trials are frequently not available and the best evidence comes from large registries with their inherent limitations of data collection and bias. It is also important to focus on intention-to-treat analysis to limit the bias associated with intolerance of therapy, which is common in this population.

References

16. de Fijter JW, Mallat MJ, Doxiadis, II, et al. Increased immunogenicity and cause of graft loss of old donor kidneys. *Journal of the American Society of Nephrology* : JASN 2001;12:1538-46.
17. Lee J, Huh KH, Park Y, et al. The clinicopathological relevance of pretransplant anti-angiotensin II type 1 receptor antibodies in renal transplantation. *Nephrology, dialysis, transplantation* : official publication of the European Dialysis and Transplant Association - European Renal Association 2015.

Guideline 3.1 – KTR: immunosuppression regimen

We recommend that the patient and/or carer should be engaged in the decisions around selection of induction agent and maintenance immunosuppression (1D)

Rationale

Immunosuppressive therapy for kidney transplantation has a wide and potentially serious side effect profile, including increased risk of cancer, cardiovascular disease and infection. Engagement with patients about anticipated short and long-term side effects of immunosuppressive therapy allows discussions to reassure about some anticipated side effects, whilst informing about others. This may aid adherence with therapy and allay untoward concerns about medication side effects.

Guideline 3.1 – KTR: Induction Immunosuppression

We recommend induction therapy should take into account the following:

- Immunosuppressive drugs should be started before or at the time of renal transplantation (1B)
- Induction therapy with biological agents should be administered to all KTRs (1B). In patients at low immunological risk this will generally involve an interleukin-2 receptor antagonist (IL2-RA). Recipients at higher immunological risk may be considered for T-cell (lymphocyte) Depleting Antibodies (TDAs; e.g. anti-lymphocyte preparations [antithymocyte globulin (ATG) or alemtuzumab])
- Induction therapy with TDAs may also be useful for lower immunological risk patients with the intention of either steroid or CNI avoidance (1C)

Guideline 3.2 – KTR: Induction Immunosuppression

We suggest that a CNI should be started at the time of transplantation and not delayed until the graft is functioning (2C)

Audit Measure

Percentage of total patients receiving induction with ILRAs and TDAs

Rationale

Following allogeneic renal transplant there is an intense period of immunological activity whereby recipient lymphocytes respond to allogeneic material. Induction therapy aims to minimise this response and the risk of early graft rejection at a time when oral agents may not have reached effective concentrations.

There is good evidence that IL2-RAs reduce the risk of early rejection when compared to placebo, although there is no definitive evidence of improved graft survival at three years, nor are there trials of adequate statistical power to answer the question of long-term benefits. There is, however, some evidence from registry data to suggest that the lower rejection rates might translate into better graft survival¹⁸. Pharmacoeconomic analysis has shown that these agents are cost effective in the early post-transplant period, and this is embodied in National Institute for Clinical Excellence (NICE) guidelines (<https://www.nice.org.uk/guidance/ta85>)¹⁹

There is moderate evidence that TDAs reduce the risk of acute rejection in high-risk immunological recipients. However, this benefit is generally gained at the expense of increased side effects - in particular, an increased incidence of malignancy, cytopenia and infection. Four randomised controlled trials comparing alemtuzumab to basiliximab in standard risk patients have consistently demonstrated reduced rates of acute rejection with alemtuzumab but longer-term outcomes are still awaited²⁰⁻²³. However, alemtuzumab has been associated with increased side effects in most studies and for most patients it would seem unnecessary other than as part of a strategy to avoid other drugs, e.g. corticosteroids. Anti-thymocyte globulin has been shown to reduce the incidence of acute rejection in standard risk KTRs when compared to an IL2RA, but follow up at ten years did not show any significant clinical differences between the two groups²⁴. A multicentre prospective RCT compared ATG to IL2RA in high risk KTRs and demonstrated significantly reduced rejection rates but no significant difference in five year outcomes²⁵. Similarly, ATG may reduce acute rejection rates in black KTRs and reduce the incidence of antibody mediated rejection (AMR) and donor-specific antibody production, but long-term evidence of benefit is lacking^{26,27}.

There is limited evidence to suggest that the clinical profile of alemtuzumab differs from that of other T cell depleting antibodies, with a lower incidence of post transplant lymphoproliferative disease (PTLD)²⁸. Preliminary evidence suggests that B lymphocyte depleting antibodies, such as anti-CD20, (rituximab) are not suitable as routine induction agents^{29,30}. Rituximab may be useful as induction agent in ABO incompatible transplantation³¹.

References

18. Lim W, Chadban S, Campbell S, Dent H, Russ G, McDonald S. Effect of interleukin-2 receptor antibody therapy on acute rejection risk and severity, long-term renal function, infection and malignancy-related mortality in renal transplant recipients. *Transplant international : official journal of the European Society for Organ Transplantation* 2010;23:1207-15.
19. Ponticelli C. Basiliximab: efficacy and safety evaluation in kidney transplantation. *Expert opinion on drug safety* 2014;13:373-81.
20. Chan K, Taube D, Roufousse C, et al. Kidney transplantation with minimized maintenance: alemtuzumab induction with tacrolimus monotherapy--an open label, randomized trial. *Transplantation* 2011;92:774-80.

21. Group CSC, Haynes R, Harden P, et al. Alemtuzumab-based induction treatment versus basiliximab-based induction treatment in kidney transplantation (the 3C Study): a randomised trial. *Lancet* 2014;384:1684-90.
22. Hanaway MJ, Woodle ES, Mulgaonkar S, et al. Alemtuzumab induction in renal transplantation. *The New England journal of medicine* 2011;364:1909-19.
23. Welberry Smith MP, Cherukuri A, Newstead CG, et al. Alemtuzumab induction in renal transplantation permits safe steroid avoidance with tacrolimus monotherapy: a randomized controlled trial. *Transplantation* 2013;96:1082-8.
24. Lentine KL, Schnitzler MA, Xiao H, Brennan DC. Long-term safety and efficacy of antithymocyte globulin induction: use of integrated national registry data to achieve ten-year follow-up of 10-10 Study participants. *Trials* 2015;16:365.
25. Hellemans R, Hazzan M, Durand D, et al. Daclizumab Versus Rabbit Antithymocyte Globulin in High-Risk Renal Transplants: Five-Year Follow-up of a Randomized Study. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2015;15:1923-32.
26. Brokhof MM, Sollinger HW, Hager DR, et al. Antithymocyte globulin is associated with a lower incidence of de novo donor-specific antibodies in moderately sensitized renal transplant recipients. *Transplantation* 2014;97:612-7.
27. Pilch NA, Taber DJ, Moussa O, et al. Prospective randomized controlled trial of rabbit antithymocyte globulin compared with IL-2 receptor antagonist induction therapy in kidney transplantation. *Annals of surgery* 2014;259:888-93.
28. Kirk AD, Cherikh WS, Ring M, et al. Dissociation of depletion induction and posttransplant lymphoproliferative disease in kidney recipients treated with alemtuzumab. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2619-25.
29. Clatworthy MR, Watson CJ, Plotnek G, et al. B-cell-depleting induction therapy and acute cellular rejection. *The New England journal of medicine* 2009;360:2683-5.
30. Macklin PS, Morris PJ, Knight SR. A systematic review of the use of rituximab as induction therapy in renal transplantation. *Transplantation reviews* 2015;29:103-8.
31. Lo P, Sharma A, Craig JC, et al. Preconditioning Therapy in ABO-Incompatible Living Kidney Transplantation: A Systematic Review and Meta-Analysis. *Transplantation* 2015.

Guideline 3.3 – KTR : Maintenance Immunosuppression

We recommend that maintenance immunosuppression should normally consist of a calcineurin inhibitor (CNI), an anti-proliferative agent with or without corticosteroids in low and medium immunological risk KTRs (1B)

Guideline 3.4 – KTR : Maintenance Immunosuppression

We suggest that low-medium dose tacrolimus (trough target 4-8ng/ml) is recommended as the CNI of choice in patients also taking steroids who are low and medium immunological risk and are not at high risk of developing post transplant diabetes mellitus (PTDM) (2C)

Guideline 3.5 – KTR : Maintenance Immunosuppression

We suggest that slow release tacrolimus preparations are used as second line agents for patients who suffer intolerable side effects related to peak dose toxicity (2C)

Guideline 3.6 – KTR : Maintenance Immunosuppression

We suggest that KTRs who are unable to tolerate tacrolimus or who suffer serious adverse reactions related to its use be considered for use of second line agents such as ciclosporin, sirolimus, everolimus, or belatacept (1B)

Guideline 3.7 – KTR : Maintenance Immunosuppression

We suggest that MPA-based drugs should be the first-line antiproliferative agent, in preference to azathioprine except in fertile KTRs who are unwilling to use reliable contraception (2B)

Guideline 3.8 – KTR : Maintenance Immunosuppression

We suggest that mycophenolate mofetil (MMF) and enteric-coated mycophenolate sodium (Myfortic[®]) provide equivalent maintenance immunosuppression (2B)

Guideline 3.9 – KTR : Maintenance Immunosuppression

We suggest that vigilant steroid avoidance or steroid withdrawal during the first week after transplantation can generally be used in low immunological risk kidney transplant recipients (2B)

Guideline 3.10 – KTR : Maintenance Immunosuppression

We suggest that minimum target levels for CNIs should be instituted in uncomplicated renal transplantation after 3 months (2C)

Guideline 3.11 – KTR : Maintenance Immunosuppression

We suggest that CNIs should be continued rather than withdrawn (2B)

Guideline 3.12 – KTR : Maintenance Immunosuppression

We suggest if steroids are not withdrawn within the first month then they should be maintained at low dose (Prednisolone - 5mg per day or less) (2C)

Audit Measures

- Percentage of *de novo* KTRs receiving tacrolimus
- Percentage of *de novo* KTRs receiving MPA-based immunosuppression
- Percentage of *de novo* KTRs receiving corticosteroid as part of maintenance therapy

Rationale

Immunosuppressive drugs are generally used in combination to balance effective total immunosuppression with minimisation of drug-specific side effects. Since the graft is most immunogenic in the early post-transplant period it is important to use higher doses of these drugs during this period. Thereafter dosages and thus blood levels can be reduced to minimise the risks of infection and malignancy. Account should be taken of the immunological risk of the transplant and also the strength of induction therapy, i.e. KTRs induced with TDAs often require less intensive maintenance therapy. High, medium and low trough (C_0) levels for tacrolimus are >10, 5-10 and <5 ng/mL respectively. Comparable C_0 levels for ciclosporin are >200, 100-200 and <100 ng/ml respectively.

It should be acknowledged that the optimal combination of immunosuppressive agents to obtain the best long-term outcomes, defined by hard endpoints such as graft and patient survival, remains elusive. There are a number of reasons for this:

- short duration of increasingly expensive trials
- convergence of outcomes with many different regimes for short term surrogate endpoints (e.g. rejection rates and graft function)
- restrictions on trial recruitment which exclude many real world patients

In the absence of such data a flexible approach is required whereby KTRs are stratified by risk and the majority are started on standard protocols. However, clinicians must be vigilant and be willing to substitute alternative agents when necessary.

Except for transplants from syngeneic or haploidentical live donors, it is generally established practice in renal transplantation to use triple therapy as maintenance, consisting of a CNI, an antiproliferative agent and corticosteroids. Such regimes have led to the lowest rejection rates and are considered the benchmark to which other regimes are compared. Induction therapy plus low-dose tacrolimus, mycophenolate mofetil (MMF) and corticosteroids have produced the lowest rates of acute rejection, superior graft function, and better graft survival. There are examples of other regimes for maintenance therapy, generally involving avoidance, minimisation or withdrawal of either steroids or CNIs. Such regimes that can produce similar outcomes in selected patients but there are currently no reliable instruments to predict which KTRs will benefit.

A large randomised controlled trial (RCT) suggested that low dose tacrolimus combined with MMF and steroids with an IL2-RA as induction was superior at 12 months in terms of graft function, graft survival and acute rejection rate to either standard or low dose ciclosporin in low immunological risk KTRs^{32,33}. There are concerns over the early nephrotoxic effects of CNIs but whether these observations extend to lower doses and levels is unknown. To date, no alternative to CNIs has been shown to improve either early or late graft outcomes. Favourable outcomes in this trial³³ were based on tacrolimus levels of 3-7 ng/mL. More recent data suggest that trough tacrolimus levels <4.0ng/mL was associated with higher levels of rejection in the 'post-SYMPHONY' era³⁴. Therefore a pragmatic compromise would be to aim for trough tacrolimus levels of 4-8ng/mL.

The risk of acute rejection is minimised by early achievement of target CNI levels and so there is no reason to delay the initiation of a CNI. Specifically there is no evidence that delaying the introduction of a CNI prevents or ameliorates delayed graft function.

Trial evidence demonstrates that tacrolimus reduces the risk of acute rejection and improves graft survival during the first year of transplantation compared to ciclosporin³⁵. Protocol biopsy studies also suggest that subclinical rejection is less prevalent in regimes containing tacrolimus as opposed to ciclosporin³⁶. However, PTDM is significantly more common with tacrolimus even accounting for variation in concomitant steroid usage³⁷. An RCT comparing tacrolimus with ciclosporin (Neoral) in non-diabetic patients demonstrated significantly higher levels of abnormalities in glucose metabolism with tacrolimus but a non-significant trend towards worse graft function with ciclosporin³⁸. However, lower blood levels of tacrolimus minimise the risk of PTDM compared to higher levels; this has not been fully explored in a trial setting.

It is acknowledged that tacrolimus is associated with a number of side effects. Some milder side effects relating to peak levels of the parent compound (e.g. tremor) may be ameliorated by dose reduction or the use of slow release formulations of tacrolimus³⁹. However, tacrolimus may cause disabling side effects in a minority of patients including posterior reversible encephalopathy syndrome, haemolytic uraemic syndrome, alopecia and GI disturbance. In such circumstances it is usually necessary to switch to a second line agent such as ciclosporin, m-TORi or belatacept.

Compared with placebo and azathioprine, mycophenolic acid based compounds, (MMF, mycophenolate sodium, generic MPA) reduces the risk of acute rejection^{40,41}. The evidence comparing MPA to placebo consistently demonstrates lower rates of acute rejection with MPA but at the expense of increased bone marrow suppression and increased opportunistic infection rates. Systemic review of the relevant studies suggests significantly reduced rejection rates and improved graft survival with MPA compared to azathioprine⁴². Absolute numbers of patients with gastrointestinal side effects are higher with MMF though this is not significant. There is limited evidence that mycophenolate sodium (Myfortic) leads to a reduced incidence of gastrointestinal side effects compared to mycophenolate mofetil, albeit in studies specifically addressing patients with gastrointestinal side effects, rather than larger studies of KTRs⁴³. All mycophenolic acid based compounds are associated with significant teratogenicity in both women and men and should not be used in fertile KTRs who are unwilling to use reliable contraception⁴⁴.

Steroids have a well-documented adverse effect profile and there is heightened interest in steroid withdrawal and avoidance regimes. Whether low dose prednisolone (e.g. 5 mg daily) is associated with a similar adverse profile to higher dose regimens is unknown. Interestingly there is little evidence of an effect on insulin sensitivity between no steroids and 5 mg daily⁴⁵. The majority of accumulated trial evidence in renal transplantation has involved steroid-containing regimes and there is little information re steroid withdrawal/avoidance. There are no differences in graft survival between patients treated with or without maintenance corticosteroids beyond the first week after kidney transplantation and avoidance beyond the first week after kidney transplantation reduces adverse effects. Early withdrawal and avoidance studies show increased acute rejection rates but without an effect on graft survival⁴⁶⁻⁴⁸. In contrast, steroid withdrawal studies later than one month after transplantation generally show increased rejection rates. Long-term follow up is required to fully assess these effects. It is clear that close vigilance is required with steroid avoidance regimes since acute rejection rates will probably be higher. Patients with graft rejection should probably be maintained on long-term oral steroids⁴⁹.

Higher doses of CNIs are required during the first three months when the recipient's immune response is receiving the most allostimulation. There is theoretically a good reason to reduce the immunosuppressive load after this time to reduce the incidence of drug-related adverse effects (i.e. reduce CNI target levels). Analysis of RCTs has shown that CNI withdrawal leads to higher rejection rates without any improvement in graft survival. Comparison of lower dose CNI regimes with higher doses have generally shown little difference in outcomes⁵⁰ but in some cases better renal function has been attained. Those seeking a fuller discussion of these studies are referred to the 2009 KDIGO guidelines⁵¹.

While there is some evidence that m-TORis can allow reduced doses of CNIs and better graft function at one year after transplantation, these agents are poorly tolerated and are associated with higher rejection rates⁵². Two studies have investigated the substitution of m-TORis for CNIs between 1 and 6 months after transplantation. While both studies demonstrated short-term improvements in renal function, there were no significant improvements in long term graft or patient outcomes⁵³⁻⁵⁵. A systemic review demonstrated a significant reduction in the rate of malignancy with sirolimus but a 43% increase in overall mortality rate compared to controls⁵⁶. The exact role of m-TORis in the early stages after transplantation is uncertain but at this time they should only be used as second line agents.

Belatacept is a fusion protein that blocks costimulatory pathways involved in T cell activation, which has to be administered by regular intravenous infusions. It has been used in KTRs as an alternative to ciclosporin and significant improvement in renal function and better graft histology at one year has been demonstrated⁵⁷. More recent studies have confirmed improved graft function, which was sustained at five years, but also an association with the development of PTLD⁵⁸. A meta-analysis concluded that belatacept has no demonstrable effect on rejection rates, patient or graft survival⁵⁹. However, it was associated with significantly better graft function and histology. It is currently a valuable second line agent that may be particularly useful in certain clinical situations (e.g. poor adherence, haemolytic uraemic syndrome). In 2011, the US Food and drug administration issued a warning over risks of PTLD and progressive multifocal leukoencephalopathy (PML) with belatacept⁶⁰.

References

32. Ekberg H, Bernasconi C, Tedesco-Silva H, et al. Calcineurin inhibitor minimization in the Symphony study: observational results 3 years after transplantation. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:1876-85.
33. Ekberg H, Tedesco-Silva H, Demirbas A, et al. Reduced exposure to calcineurin inhibitors in renal transplantation. *The New England journal of medicine* 2007;357:2562-75.
34. Gaynor JJ, Ciancio G, Guerra G, et al. Lower tacrolimus trough levels are associated with subsequently higher acute rejection risk during the first 12 months after kidney transplantation. *Transplant international : official journal of the European Society for Organ Transplantation* 2016;29:216-26.
35. Webster AC, Woodroffe RC, Taylor RS, Chapman JR, Craig JC. Tacrolimus versus ciclosporin as primary immunosuppression for kidney transplant recipients: meta-analysis and meta-regression of randomised trial data. *Bmj* 2005;331:810.
36. Rowshani AT, Scholten EM, Bemelman F, et al. No difference in degree of interstitial Sirius red-stained area in serial biopsies from area under concentration-over-time curves-guided cyclosporine versus tacrolimus-treated renal transplant recipients at one year. *Journal of the American Society of Nephrology : JASN* 2006;17:305-12.
37. Woodward RS, Schnitzler MA, Baty J, et al. Incidence and cost of new onset diabetes mellitus among U.S. wait-listed and transplanted renal allograft recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2003;3:590-8.
38. Vincenti F, Friman S, Scheuermann E, et al. Results of an international, randomized trial comparing glucose metabolism disorders and outcome with cyclosporine versus tacrolimus. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:1506-14.
39. Langone A, Steinberg SM, Gedaly R, et al. Switching STudy of Kidney TRansplant PATients with Tremor to LCP-TacrO (STRATO): an open-label, multicenter, prospective phase 3b study. *Clinical transplantation* 2015;29:796-805.
40. Mycophenolate Mofetil Acute Renal Rejection Study G. Mycophenolate mofetil for the treatment of a first acute renal allograft rejection: three-year follow-up. The Mycophenolate Mofetil Acute Renal Rejection Study Group. *Transplantation* 2001;71:1091-7.
41. Shapiro R, Jordan ML, Scantlebury VP, et al. A prospective, randomized trial of tacrolimus/prednisone vs tacrolimus/prednisone/mycophenolate mofetil in renal transplantation: 1-year actuarial follow-up. *Transplantation proceedings* 1999;31:1134.
42. Wagner M, Earley AK, Webster AC, Schmid CH, Balk EM, Uhlig K. Mycophenolic acid versus azathioprine as primary immunosuppression for kidney transplant recipients. *The Cochrane database of systematic reviews* 2015;12:CD007746.
43. Shehata M, Bhandari S, Venkat-Raman G, et al. Effect of conversion from mycophenolate mofetil to enteric-coated mycophenolate sodium on maximum tolerated dose and gastrointestinal symptoms following kidney transplantation. *Transplant international : official journal of the European Society for Organ Transplantation* 2009;22:821-30.
44. Mycophenolate mofetil, mycophenolic acid: new pregnancy-prevention advice for women and men. 2015. at <https://www.gov.uk/drug-safety-update/mycophenolate-mofetil-mycophenolic-acid-new-pregnancy-prevention-advice-for-women-and-men>.)
45. Midtvedt K, Hjelmesaeth J, Hartmann A, et al. Insulin resistance after renal transplantation: the effect of steroid dose reduction and withdrawal. *Journal of the American Society of Nephrology : JASN* 2004;15:3233-9.
46. Matas AJ, Gillingham K, Kandaswamy R, et al. Kidney transplant half-life (t[1/2]) after rapid discontinuation of prednisone. *Transplantation* 2009;87:100-2.
47. Matas AJ, Kandaswamy R, Gillingham KJ, et al. Prednisone-free maintenance immunosuppression-a 5-year experience. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2005;5:2473-8.
48. Woodle ES, First MR, Pirsch J, Shihab F, Gaber AO, Van Veldhuisen P. A prospective, randomized, double-blind, placebo-controlled multicenter trial comparing early (7 day) corticosteroid cessation versus long-term, low-dose corticosteroid therapy. *Annals of surgery* 2008;248:564-77.

49. Humar A, Gillingham K, Kandaswamy R, Payne W, Matas A. Steroid avoidance regimens: a comparison of outcomes with maintenance steroids versus continued steroid avoidance in recipients having an acute rejection episode. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:1948-53.
50. Knight SR, Morris PJ. Does the evidence support the use of mycophenolate mofetil therapeutic drug monitoring in clinical practice? A systematic review. *Transplantation* 2008;85:1675-85.
51. KDIGO clinical practice guideline for the care of kidney transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9 Suppl 3:S1-155.
52. Webster AC, Lee VW, Chapman JR, Craig JC. Target of rapamycin inhibitors (sirolimus and everolimus) for primary immunosuppression of kidney transplant recipients: a systematic review and meta-analysis of randomized trials. *Transplantation* 2006;81:1234-48.
53. Budde K, Becker T, Arns W, et al. Everolimus-based, calcineurin-inhibitor-free regimen in recipients of de-novo kidney transplants: an open-label, randomised, controlled trial. *Lancet* 2011;377:837-47. doi: 10.1016/S0140-6736(10)62318-5. Epub 2011 Feb 19.
54. Budde K, Lehner F, Sommerer C, et al. Five-year outcomes in kidney transplant patients converted from cyclosporine to everolimus: the randomized ZEUS study. *Am J Transplant* 2015;15:119-28. doi: 10.1111/ajt.12952. Epub 2014 Dec 17.
55. Weir MR, Mulgaonkar S, Chan L, et al. Mycophenolate mofetil-based immunosuppression with sirolimus in renal transplantation: a randomized, controlled Spare-the-Nephron trial. *Kidney Int* 2011;79:897-907. doi: 10.1038/ki.2010.492. Epub Dec 29.
56. Knoll GA, Kokolo MB, Mallick R, et al. Effect of sirolimus on malignancy and survival after kidney transplantation: systematic review and meta-analysis of individual patient data. *BMJ* 2014;349:g6679. doi:10.1136/bmj.g6679.
57. Vincenti F, Larsen C, Durrbach A, et al. Costimulation blockade with belatacept in renal transplantation. *N Engl J Med* 2005;353:770-81.
58. Rostaing L, Vincenti F, Grinyo J, et al. Long-term belatacept exposure maintains efficacy and safety at 5 years: results from the long-term extension of the BENEFIT study. *Am J Transplant* 2013;13:2875-83. doi: 10.1111/ajt.12460. Epub 2013 Sep 18.
59. Masson P, Henderson L, Chapman JR, Craig JC, Webster AC. Belatacept for kidney transplant recipients. *Cochrane Database Syst Rev* 2014;11:CD010699. doi:10.1002/14651858.CD010699.pub2.
60. Nulojix (belatacept): Risk Evaluation and Mitigation Strategy (REMS). 2011. at <http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm262210.htm>.)

Guideline 3.13 – KTR : Monitoring of Immunosuppression

We suggest that long-term monitoring of immunosuppression levels is required as follows:

- Tacrolimus and ciclosporin levels should be monitored. The frequency should be three times a week immediately after the transplant. Levels should be checked when any medication with possible interactions is prescribed or when there is unexplained graft dysfunction (2C)
- Tacrolimus should be monitored by the C₀ trough level, while ciclosporin can be monitored by either trough (C₀) or 2-hour post dose (C₂) level (2C)
- Tacrolimus and ciclosporin levels should be available within 24 hours of taking blood samples (2C)
- The utility of monitoring mycophenolic acid (MPA) C₀ levels is uncertain (2D)
- Sirolimus should be monitored by the C₀ trough level (2C)

Rationale

Therapeutic drug monitoring is advisable for drugs with a narrow therapeutic index. For tacrolimus and ciclosporin the absorption may vary in the early stages after transplantation but usually stabilises within a month. Both drugs may exhibit both inter-patient and intra-patient variability. Tacrolimus and ciclosporin are traditionally monitored by 12-hour C_0 trough levels, but in the case of ciclosporin there is some evidence that C_2 levels may also be used although target ranges are less well established and the logistics of sample collection are more complex. There is little evidence directly comparing different target levels of the same drug in a controlled fashion. Drug monitoring of MMF is best carried out by measuring the area under the curve (AUC) but clinical studies have not been conclusive^{50,61,62}. C_0 levels correlate poorly with AUC and remain unproven and rarely used in clinical practice. Sirolimus levels should be monitored since toxic effects correlate with high drug levels and C_0 levels correlate well with AUC^{63,64}.

References

61. Kuypers DR, Bammens B, Claes K, Evenepoel P, Vanrenterghem Y. Maintenance immunosuppressive agents as risk factors for BK virus nephropathy: the need for true drug exposure measurements. *Transplantation* 2010;89:1296-7; author reply 7-8.
62. van Gelder T. Mycophenolate blood level monitoring: recent progress. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:1495-9.
63. Kahan BD, Camardo JS. Rapamycin: clinical results and future opportunities. *Transplantation* 2001;72:1181-93.
64. Kovarik JM, Tedesco H, Pascual J, et al. Everolimus therapeutic concentration range defined from a prospective trial with reduced-exposure cyclosporine in de novo kidney transplantation. *Ther Drug Monit* 2004;26:499-505.

Guideline 3.14 – KTR : Prescribing and the use of Generic agents

We suggest that generic immunosuppression compounds should not be used unless they have been shown to be bioequivalent and approved by the European Agency for the Evaluation of Medicinal Products (2D)

Guideline 3.15 – KTR : Prescribing and the use of Generic agents

We suggest that KTRs should be made aware of the existence of generics and the dangers of indiscriminate usage (2D)

Guideline 3.16 – KTR : Prescribing and the use of Generic agents

We suggest that drugs should be prescribed by brand name where unproven generic substitutes are available (2D)

Guideline 3.17 – KTR : Prescribing and the use of Generic agents

We suggest that KTRs should be followed closely after switching to a generic preparation until a new steady state is established (2D)

Audit Measure

- The use of generic agents should be monitored and audited

Rationale

The introduction of many generic preparations of tacrolimus, ciclosporin and MPA has occurred in the last five years. These offer potential cost savings but with the risk that these medications are not truly bioequivalent due to the limitations of the regulatory process. This has led to differences in both pharmacokinetic and clinical terms, compared with the original agents.

Immunosuppressive drugs in common use have narrow therapeutic windows, with significant risk of under- and over-immunosuppression and, with CNIs, the risk of nephrotoxicity due to over-exposure. Plasma CNI levels are carefully measured in practice, with narrow therapeutic ranges. Assessment of generic agents requires only that time-averaged plasma concentrations (area-under-the-curve) fall between 80-125% of the original preparation in normal subjects. Differences in bioavailability due to food, or other factors, are not assessed. For ciclosporin the bioavailability of generic agents extends across this range and is influenced by food, thus switching between generic agents may result in major differences in drug exposure. This may be minimised by careful measure of drug exposure after switching between agents and generic preparations (“named” generics). When the choice of generic is left to the dispenser this is likely to result in variable exposure. The same issues may apply to tacrolimus but the bioavailability of this agent is less variable, and to other generic immunosuppressants such as MPA, although monitoring of this agent is not usually undertaken in clinical practice. For these reasons a local protocol for the use of generic agents should be available to all involved in the care of transplant recipients, specifically those who write and dispense prescriptions.

4. Kidney Transplant Recipient (KTR): Acute rejection (Guidelines 4.1-4.12)

Guideline 4.1 – KTR: Diagnosis of acute rejection

We recommend that a transplant renal biopsy should be carried out before treating an acute rejection episode unless this will substantially delay treatment or pose a significant risk to the patient (1C)

Guideline 4.2 – KTR: Diagnosis of acute rejection

We suggest that two cores of renal tissue should be obtained if possible since this will increase the sensitivity of the investigation (2C)

Guideline 4.3 – KTR: Diagnosis of acute rejection

We recommend that a 16 gauge automated core biopsy needle is used where possible to provide the best compromise between diagnostic usefulness and patient tolerance of the procedure (1C)

Guideline 4.4 – KTR: Diagnosis of acute rejection

We recommend that a protocol transplant renal biopsy, defined as a biopsy performed in a stable graft without clinical evidence of acute rejection (proteinuria, rising creatinine), be considered in the setting of persisting delayed graft function (DGF) (1C)

Guideline 4.5 – KTR: Diagnosis of acute rejection

We recommend that routine C4d and SV40 staining should be performed upon transplant biopsies (2C)

Guideline 4.6 – KTR: Diagnosis of acute rejection

We suggest that a serum sample be sent at the time of renal biopsy (for graft dysfunction) to look for HLA-specific antibodies (2C)

Audit measures

- Severity of biopsy proven acute rejection (BPAR) recorded by Banff criteria
- The percentage of KTRs with BPAR in first 3 months and the first 12 months
- The percentage of KTRs requiring TDAs to treat rejection within the first year
- Complication rates after renal transplant biopsy
- The percentage of KTRs with a donor-specific HLA antibody at the time of biopsy

Rationale

Historically, unresolved acute rejection episodes invariably led to graft loss so it is rational to treat such episodes unless the treatment is likely to do more harm than good. Rejection episodes are characteristically associated with loss of graft function but diagnosis is best established by a percutaneous biopsy since it differentiates rejection clearly from other causes of graft dysfunction. Recognition of different forms of rejection may inform different treatment regimens (e.g. antibody mediated rejection).

Two cores of tissue should be obtained as this approach establishes the diagnosis of acute rejection with a sensitivity of 99% versus 91% when only one core is obtained⁶⁵. The use of a 16 gauge automated core biopsy needle yields higher numbers of glomeruli and thus increased diagnostic usefulness without an increase in complication rate compared with an 18 gauge needle^{66,67}.

Biopsy is performed when there is acute graft dysfunction with the aim of providing a histological diagnosis to confirm clinical suspicion. However, renal scarring may be too advanced for any intervention to lead to significant improvement. Subclinical acute rejection⁶⁸ is defined as histological rejection in the absence of clinical evidence of altered graft function diagnosed on protocol biopsies and logically one might expect that early detection would lead to improved outcomes. In most patients, with modern immunosuppression regimes, there is little evidence that treatment of SCAR improves outcomes and a clear link between SCAR and chronic rejection has not yet been proven⁶⁹. There are multiple reasons for this: there is no large multi-centre prospective study; the induction, baseline and maintenance immunosuppression regimes vary in studies; some protocol biopsies are actually indication biopsies and thus cannot show SCAR; and the Banff criteria was designed for diagnostic biopsies and not for protocol biopsies where the findings may not fit with the criteria⁷⁰. There is randomised control trial evidence that there is no benefit from protocol biopsies performed within the first 6 months of transplantation in low risk patients on a standard immunosuppressive regimen⁶⁹. There are two clinical settings where protocol biopsy is of value. In delayed graft function (DGF) there is increased risk of graft failure; 'silent' acute rejection may account for a significant proportion of this increased risk⁷¹. In a more specialised setting, in very high and high immunological risk patients, if SCAR is detected this should be treated as it may improve outcome^{72,73}. The optimal timing and frequency of protocol biopsy is not clear but we suggest that it be considered in the clinical settings outlined above at days 7-10 for DGF. In high-risk transplants, protocol biopsy will typically be at 3 months.

It is recommended to stain all biopsies for C4d and to send a serum sample for detection of human leukocyte antigen (HLA) specific antibodies to facilitate diagnosis of acute antibody mediated rejection in line with joint BSHI/BTS guidelines and those of The Transplantation Society^{73,74}. The Banff criteria were revised in 2013 to acknowledge the occurrence of antibody mediated rejection in the absence of detectable C4d staining⁷⁰. Immunohistopathologic evidence of recent interaction of antibody with the vascular endothelium is necessary for the diagnosis and this may include, but is not limited, to C4d positivity. Conversely, diffuse C4d staining may occur in the absence of morphological evidence of active rejection or graft dysfunction, primarily in ABO incompatible transplantation.

An episode of acute rejection is a period of uncertainty and is likely to cause anxiety in KTR and/or their carers. Good communication explaining the treatment rationale and likely outcomes of rejection is important in addressing these concerns.

References

65. Colvin RB, Cohen AH, Saiontz C, et al. Evaluation of pathologic criteria for acute renal allograft rejection: reproducibility, sensitivity, and clinical correlation. *Journal of the American Society of Nephrology : JASN* 1997;8:1930-41.
66. Nicholson ML, Wheatley TJ, Doughman TM, et al. A prospective randomized trial of three different sizes of core-cutting needle for renal transplant biopsy. *Kidney Int* 2000;58:390-5.
67. Peters B, Andersson Y, Stegmayr B, et al. A study of clinical complications and risk factors in 1,001 native and transplant kidney biopsies in Sweden. *Acta radiologica* 2014;55:890-6.
68. Schena FP, Pascoe MD, Alberu J, et al. Conversion from calcineurin inhibitors to sirolimus maintenance therapy in renal allograft recipients: 24-month efficacy and safety results from the CONVERT trial. *Transplantation* 2009;87:233-42.
69. Rush D, Arlen D, Boucher A, et al. Lack of benefit of early protocol biopsies in renal transplant patients receiving TAC and MMF: a randomized study. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2538-45.
70. Haas M, Sis B, Racusen LC, et al. Banff 2013 meeting report: inclusion of c4d-negative antibody-mediated rejection and antibody-associated arterial lesions. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2014;14:272-83.
71. Qureshi F, Rabb H, Kasiske BL. Silent acute rejection during prolonged delayed graft function reduces kidney allograft survival. *Transplantation* 2002;74:1400-4.
72. Kraus ES, Parekh RS, Oberai P, et al. Subclinical rejection in stable positive crossmatch kidney transplant patients: incidence and correlations. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:1826-34.
73. Tait BD, Susal C, Gebel HM, et al. Consensus guidelines on the testing and clinical management issues associated with HLA and non-HLA antibodies in transplantation. *Transplantation* 2013;95:19-47.
74. BSHI/BTS. Guidelines for the detection and characterisation of clinically relevant antibodies in allotransplantation. 2014.

Guideline 4.7 – KTR: Treatment of acute rejection

We suggest that borderline acute cellular rejection should be treated in the context of acute graft dysfunction (2D)

Guideline 4.8 – KTR: Treatment of acute rejection

We recommend that high dose intravenous corticosteroids should be the first line treatment for acute cellular rejection (1D)

Guideline 4.9 – KTR: Treatment of acute rejection

We suggest that maintenance steroids should be added or restarted in steroid-free patients undergoing acute rejection of any type (2D)

Guideline 4.10 – KTR: Treatment of acute rejection

We suggest that lymphocyte depleting agents may be considered for refractory acute cellular rejection or aggressive vascular cellular rejection (i.e. Banff category 4 Type II and III) (2C)

Guideline 4.11.1 – KTR: Treatment of acute rejection

We suggest that antibody mediated rejection (AMR) should be treated with one or more of the following modalities: steroids; plasma exchange; intravenous immunoglobulin; anti-CD20 antibody; lymphocyte-depleting antibody or bortezomib (2C)

Guideline 4.11.2 - KTR: Treatment of acute rejection

We recommend that the BTS guidelines on antibody incompatible transplantation for management of rejection in the context of antibody incompatible transplantation (1A-D)

Guideline 4.12 – KTR: Treatment of acute rejection

We suggest after an episode of rejection (unless associated with low CNI levels) that azathioprine should be switched to MPA-based immunosuppression, MPA should be started or the existing dose of MPA maximised and ciclosporin and sirolimus should be switched to tacrolimus (2D)

Rationale

The management of antibody incompatible transplantation is reviewed extensively in the recent BTS guidelines on this topic⁷⁵. In contrast to SCAR, borderline acute cellular rejection detected in the context of graft dysfunction should be treated in the knowledge that if untreated, the infiltrates may progress into rejection with consequent deterioration in transplant function^{76,77}. However, there is little evidence to guide therapy, treatment is controversial, and there is evidence that borderline infiltrates will not automatically progress into rejection^{77,78}.

The majority of acute cellular rejection episodes respond to treatment with corticosteroids^{79,80}. The optimal regime for steroid administration has not been determined but intravenous methylprednisolone on three consecutive days is commonly used⁷⁹. If intravenous steroid is precluded, high dose oral steroid can be utilised. The use of T cell depleting antibodies (TDAs) in milder grades of cellular rejection (Banff category 4 type I) may be more effective in restoring renal function but results in significantly greater side effects^{81,82}. If renal function does not return to baseline with steroid and/or ATG, or if there is a new decline in function after successful treatment of an acute rejection episode, a repeat biopsy should be considered to rule out additional rejection or other causes of graft dysfunction (e.g. concurrent acute tubular necrosis or BK nephropathy). Treating more severe cellular rejection (Banff category 4 Type IIa, IIb or III) and steroid unresponsive episodes with TDAs often improves graft function although a thorough risk-benefit assessment of such treatment should be undertaken^{81,82}. There is some evidence that adding an MPA product after such episodes or substituting azathioprine with MPA will result in fewer subsequent rejection episodes⁴⁰. Intensifying immunosuppression after a rejection episode may help prevent further rejection including the following: maximising the dose of MPA product and switching ciclosporin or sirolimus to tacrolimus⁸³⁻⁸⁶.

If AMR is diagnosed, there is limited evidence that treatment with alternative modalities including plasmapheresis, immunoabsorption, intravenous immunoglobulin or monoclonal antibodies may be beneficial⁸⁷. Trial evidence is conflicting and of low quality. Small non-randomised studies and case reports suggest that monoclonal antibodies targeting B cell function and thus antibody production (rituximab and bortezomib) may be of benefit, as may terminal complement inhibition with eculizumab. However, there is insufficient evidence to recommend these agents and the risks and benefits, particularly of biologic therapies, must be considered on an individual basis⁸⁷⁻⁸⁹.

References

75. BTS. Antibody Incompatible Transplantation Guideline. 2015.
76. Matoza JR, Danguilan RA, Chicano S. Impact of Banff borderline acute rejection among renal allograft recipients. *Transplantation proceedings* 2008;40:2303-6.
77. Wu K, Budde K, Lu H, et al. The severity of acute cellular rejection defined by Banff classification is associated with kidney allograft outcomes. *Transplantation* 2014;97:1146-54.
78. Beimler J, Zeier M. Borderline rejection after renal transplantation--to treat or not to treat. *Clinical transplantation* 2009;23 Suppl 21:19-25.
79. Gray D, Shepherd H, Daar A, Oliver DO, Morris PJ. Oral versus intravenous high-dose steroid treatment of renal allograft rejection. The big shot or not? *Lancet* 1978;1:117-8.

80. Shinn C, Malhotra D, Chan L, Cosby RL, Shapiro JI. Time course of response to pulse methylprednisolone therapy in renal transplant recipients with acute allograft rejection. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 1999;34:304-7.
81. Solez K, Colvin RB, Racusen LC, et al. Banff 07 classification of renal allograft pathology: updates and future directions. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2008;8:753-60.
82. Webster AC, Pankhurst T, Rinaldi F, Chapman JR, Craig JC. Monoclonal and polyclonal antibody therapy for treating acute rejection in kidney transplant recipients: a systematic review of randomized trial data. *Transplantation* 2006;81:953-65.
83. Chhabra D, Skaro AI, Leventhal JR, et al. Long-term kidney allograft function and survival in prednisone-free regimens: tacrolimus/mycophenolate mofetil versus tacrolimus/sirolimus. *Clinical journal of the American Society of Nephrology : CJASN* 2012;7:504-12.
84. Morrissey PE, Gohh R, Shaffer D, et al. Correlation of clinical outcomes after tacrolimus conversion for resistant kidney rejection or cyclosporine toxicity with pathologic staging by the Banff criteria. *Transplantation* 1997;63:845-8.
85. Sun Q, Liu ZH, Cheng Z, et al. Treatment of early mixed cellular and humoral renal allograft rejection with tacrolimus and mycophenolate mofetil. *Kidney Int* 2007;71:24-30.
86. Sun Q, Liu ZH, Yin G, et al. Tacrolimus combined with mycophenolate mofetil can effectively reverse C4d-positive steroid-resistant acute rejection in Chinese renal allograft recipients. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2006;21:510-7.
87. Roberts DM, Jiang SH, Chadban SJ. The treatment of acute antibody-mediated rejection in kidney transplant recipients-a systematic review. *Transplantation* 2012;94:775-83.
88. Lefaucheur C, Nochy D, Andrade J, et al. Comparison of combination Plasmapheresis/IVIg/anti-CD20 versus high-dose IVIg in the treatment of antibody-mediated rejection. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:1099-107.
89. Locke JE, Magro CM, Singer AL, et al. The use of antibody to complement protein C5 for salvage treatment of severe antibody-mediated rejection. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:231-5.

5. Kidney Transplant Recipient (KTR): Chronic Allograft Injury (CAI) Guidelines 5.1-5.)

Guideline 5.1 – KTR: Diagnosis of Chronic Allograft Injury (CAI)

We recommend that early identification of graft injury is desirable to maximise the potential to intervene. A proactive and systematic approach should be employed to manage graft dysfunction (1C)

Guideline 5.2 – KTR: Detection of Chronic Allograft Injury (CAI)

We suggest that renal function should be monitored at each clinic visit by assessment of serum creatinine and qualitative evaluation of urine protein excretion by dipstick supplemented by spot PCR or ACR if positive (2C)

Guideline 5.3 – KTR: Diagnosis of Chronic Allograft Injury (CAI)

We suggest that renal biopsy is the optimal investigation for parenchymal causes of graft dysfunction where the cause is uncertain (2C)

Guideline 5.4 – KTR: Diagnosis of Chronic Allograft Injury (CAI)

We suggest that renal biopsies in patients with chronically deteriorating function should routinely be stained for C4d and SV40 (2C)

Guideline 5.5 – KTR: Diagnosis of Chronic Allograft Injury (CAI)

We suggest that a serum sample should be sent at the time of renal biopsy (for graft dysfunction) to look for HLA-specific antibodies (2C)

Audit Measures

- Severity of CAI recorded by BANFF criteria
- The percentage of KTRs with positive C4d staining on biopsy
- The percentage of KTRs with a donor specific HLA antibody at the time of biopsy

Rationale

Unfortunately there are currently no good markers of early allograft injury. A number of non-invasive biomarkers have been proposed including urine and serum microRNA profiling but the clinical utility and additive predictive value of these compared to traditional markers – serum creatinine, proteinuria, histopathology – is uncertain⁹⁰⁻⁹⁴. Graft damage can be detected by protocol biopsy but the utility of this approach is unproven. Studies show that all protocol biopsies at 3 years post transplantation display evidence of some degree of CAI and by 5 years this is classed as moderate or severe in over 60% of patients⁹⁵. Therefore current best practice consists of vigilant monitoring of simple clinical markers of allograft function including serum creatinine and proteinuria^{96,97}. More complex and expensive approaches such as monitoring serum anti-HLA antibodies also remain unproven.

Deterioration in graft function is a heterogeneous entity with multiple causes, both immunological and non-immunological^{98,99}. Treatment may entail diametrically opposite strategies and therefore deterioration of allograft function should be investigated by percutaneous biopsy if possible. Tissue samples should be examined by an experienced renal histopathologist and classified according to the Banff criteria⁷⁰. Staining for C4d deposition and SV40 antigen should be routinely available because positive staining will affect the treatment strategy.

Although there is not yet any proven therapy, it is important to recognise chronic humoral rejection diagnosed according to the Banff criteria^{70,81,100}. The detection of anti-HLA antibodies and C4d staining on transplant biopsy are associated with worse clinical outcomes^{99,101-107}. More recently the complement binding ability of anti-HLA antibodies has been shown to impact upon graft survival¹⁰⁸. Post-transplant screening for anti-HLA antibodies has been suggested but there is not yet a solid evidence base to recommend it routinely in low or moderate immunological risk patients, nor to routinely determine the complement fixing ability of detected anti-HLA antibodies. In patients who are sensitised or deemed high immunological risk, a timetable of post-transplant antibody sampling should be agreed within the transplant centre. This is in line with the recommendations set out in both the BSHI/BTS and the Transplantation Society Guidelines^{73,74}.

Identifying a process likely to lead to a decline in long term transplant function will be worrying for patients/their carers. It is important to involve patients in treatment decisions including need for biopsy, alterations to the immunosuppressive regime. This may also include plans for return to dialysis or re-transplantation.

References

90. Amer H, Fidler ME, Myslak M, et al. Proteinuria after kidney transplantation, relationship to allograft histology and survival. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2748-56.
91. Amer H, Lieske JC, Rule AD, et al. Urine high and low molecular weight proteins one-year post-kidney transplant: relationship to histology and graft survival. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2013;13:676-84.

92. Hartono C, Muthukumar T, Suthanthiran M. Noninvasive diagnosis of acute rejection of renal allografts. *Current opinion in organ transplantation* 2010;15:35-41.
93. Ho J, Wiebe C, Gibson IW, Rush DN, Nickerson PW. Immune monitoring of kidney allografts. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2012;60:629-40.
94. Maluf DG, Dumur CI, Suh JL, et al. The urine microRNA profile may help monitor post-transplant renal graft function. *Kidney Int* 2014;85:439-49.
95. Nankivell BJ, Borrows RJ, Fung CL, O'Connell PJ, Allen RD, Chapman JR. Natural history, risk factors, and impact of subclinical rejection in kidney transplantation. *Transplantation* 2004;78:242-9.
96. Amer H, Cosio FG. Significance and management of proteinuria in kidney transplant recipients. *Journal of the American Society of Nephrology : JASN* 2009;20:2490-2.
97. Grinyo JM, Saval N, Campistol JM, Group IS. Clinical assessment and determinants of chronic allograft nephropathy in maintenance renal transplant patients. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2011;26:3750-5.
98. El-Zoghby ZM, Stegall MD, Lager DJ, et al. Identifying specific causes of kidney allograft loss. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:527-35.
99. Sellares J, de Freitas DG, Mengel M, et al. Understanding the causes of kidney transplant failure: the dominant role of antibody-mediated rejection and nonadherence. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2012;12:388-99.
100. Moll S, Pascual M. Humoral rejection of organ allografts. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2005;5:2611-8.
101. Bohmig GA, Exner M, Habicht A, et al. Capillary C4d deposition in kidney allografts: a specific marker of alloantibody-dependent graft injury. *Journal of the American Society of Nephrology : JASN* 2002;13:1091-9.
102. Everly MJ, Rebellato LM, Haisch CE, et al. Incidence and impact of de novo donor-specific alloantibody in primary renal allografts. *Transplantation* 2013;95:410-7.
103. Hidalgo LG, Campbell PM, Sis B, et al. De novo donor-specific antibody at the time of kidney transplant biopsy associates with microvascular pathology and late graft failure. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:2532-41.
104. Kedainis RL, Koch MJ, Brennan DC, Liapis H. Focal C4d+ in renal allografts is associated with the presence of donor-specific antibodies and decreased allograft survival. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:812-9.
105. Lachmann N, Terasaki PI, Budde K, et al. Anti-human leukocyte antigen and donor-specific antibodies detected by luminex posttransplant serve as biomarkers for chronic rejection of renal allografts. *Transplantation* 2009;87:1505-13.
106. Mao Q, Terasaki PI, Cai J, et al. Extremely high association between appearance of HLA antibodies and failure of kidney grafts in a five-year longitudinal study. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:864-71.
107. Wiebe C, Gibson IW, Blydt-Hansen TD, et al. Evolution and clinical pathologic correlations of de novo donor-specific HLA antibody post kidney transplant. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2012;12:1157-67.
108. Loupy A, Lefaucheur C, Vernerey D, et al. Complement-binding anti-HLA antibodies and kidney-allograft survival. *The New England journal of medicine* 2013;369:1215-26.

Guideline 5.6 – KTR: Treatment of chronic allograft injury

We suggest that chronic allograft injury should be treated:

- By withdrawal of calcineurin inhibitors (CNIs) if there is either histological evidence of CNI toxicity or non-specific interstitial fibrosis and tubular atrophy (2C)
- By intensification of immunosuppression if there is evidence of ongoing immune injury (cellular rejection and/or humoral rejection) (2C)

In a similar fashion to other patients with CKD following similar preventative strategies and with timely referral to low clearance services (2D)

Rationale

There is some evidence that withdrawal of CNIs following chronic deterioration of graft function is beneficial^{55,109,110}. The role of m-TOR-inhibitors as replacements for CNIs is uncertain but this approach should be avoided in patients with eGFR <40 mL/min/1.73m² and/or significant proteinuria (PCR >50 or ACR >35 mg/mmol)⁶⁸. An overview of treatment options dictated by findings on allograft biopsy is shown in Table 2.

Banff code	Descriptive term	Pathophysiology	Interpretation	Treatment
i	Interstitial inflammation	Infiltration of interstitium by mononuclear cells	Linked with cellular rejection but also viral infection	Intensification of immunosuppression –often pulsed intravenous steroids
t	Tubulitis	Infiltration of renal tubules by mononuclear cells	Linked with cellular rejection but also viral infection	Intensification of immunosuppression –often pulsed intravenous steroids
g	Glomerulitis	Margination of inflammatory leukocytes in the glomerular capillary loops	Marker of humeral rejection	Intensification of immunosuppression if not too much chronic damage
v	Arterial inflammation	Inflammation of arterial wall with infiltration of mononuclear cells	Marker of either severe cellular rejection or humeral rejection	Intensification of immunosuppression if not too much chronic damage
ptc	Peritubular capillaritis	Margination of inflammatory cells in the peritubular capillaries	Marker of humeral rejection	Intensification of immunosuppression if not too much chronic damage
ci	Interstitial Fibrosis	Interstitial structure replaced by fibrosis	Marker of chronic damage	Poor prognostic sign – may prompt reduction in CNI
ct	Tubular atrophy	Interstitial tubules involuted	Marker of chronic damage	Poor prognostic sign – may prompt reduction in CNI
cg	Transplant glomerulopathy	Interposition of mesangium and thickening of GBM	Associated with proteinuria and development of DSAs – End lesion of CAMR	Poor prognosis – no known treatment but intensification of immunosuppression often practiced
mm	Mesangial matrix expansion	Increase of thickness of mesangial matrix	Marker of microvascular damage to glomerulus	Usually interpreted in association with other findings
cv	Arterial fibrointimal thickening	Expansion of intima between endothelium and media	Marker of chronic damage – non-specific	Poor prognostic sign – vascular protective measures
ah	Arteriolar hyalinosis	Nodular deposition of hyaline	CNI toxicity but non-specific (e.g. HT, DM, lipids)	Reduction or withdrawal of CNI

Table 2 – Interpretation of the Banff Classification

There is no proven therapy for chronic humoral rejection and studies are ongoing in this area. Whilst to date there is limited high quality (e.g. randomised controlled trials) evidence to support this strategy, it seems logical to consider increased immunosuppression in the face of ongoing immunological damage to the kidney transplant. Employing measures used in other non-transplant patients with CKD is likely to be of benefit; for example, some studies show that anaemia is more prevalent in transplant patients and is associated with poor outcomes¹¹¹. The BTS 'Management of the Failing Kidney Transplant' guideline covers this aspect of management in detail¹¹².

References

109. Dudley C, Pohanka E, Riad H, et al. Mycophenolate mofetil substitution for cyclosporine a in renal transplant recipients with chronic progressive allograft dysfunction: the "creeping creatinine" study. *Transplantation* 2005;79:466-75.
110. Pascual M, Theruvath T, Kawai T, Tolkoff-Rubin N, Cosimi AB. Strategies to improve long-term outcomes after renal transplantation. *The New England journal of medicine* 2002;346:580-90.
111. Winkelmayr WC, Chandraker A. Posttransplantation anemia: management and rationale. *Clinical journal of the American Society of Nephrology : CJASN* 2008;3 Suppl 2:S49-55.
112. BTS. Management of the Failing Kidney Transplant. 2013.

Guideline 5.7 – KTR: Renal biopsy in chronic allograft injury

We suggest that a renal transplant biopsy is indicated:

- If there is a persistent unexplained elevation of creatinine or failure to return to baseline after an episode of biopsy proven acute rejection (BPAR) (1C)
- Every 7-10 days during delayed graft function (DGF) (2C)
- If expected renal function is not achieved within 4-8 weeks (2D)
- If sustained new onset proteinuria develops (PCR >50 mg/mmol or ACR >35 mg/mmol) (2C)

Audit Measures

- The percentage of KTRs with positive C4d staining on biopsy
- The percentage of KTRs with a donor specific HLA antibody at the time of biopsy

Rationale

Unexplained changes in serum creatinine are an important clinical marker of rejection and proteinuria is associated with poor outcome; both should be investigated for a treatable cause^{90,113,114}.

References

113. Cherukuri A, Welberry-Smith MP, Tattersall JE, et al. The clinical significance of early proteinuria after renal transplantation. *Transplantation* 2010;89:200-7.
114. Halimi JM, Laouad I, Buchler M, et al. Early low-grade proteinuria: causes, short-term evolution and long-term consequences in renal transplantation. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2005;5:2281-8.

6. Kidney Transplant Recipient (KTR): Cardiovascular disease and lifestyle (Guidelines 6.1-6.6)

Guideline 6.1– KTR: Hypertension

We suggest that the management of hypertension take into account that:

- Blood pressure should be recorded at each clinic visit (1C)
- Clinic blood pressure should be <140/90 mmHg in clinic (130/80 mmHg if PCR >50 or ACR >35) (2C)
- Home blood pressure recordings and 24-hour ambulatory recordings may be helpful in some instances but lower BP targets should then be set (home and or ambulatory daytime measures <135/80mmHg) (2D)
- There is no evidence that any antihypertensive agent is better than any other and effort should be focused on achieving absolute levels rather than the use of individual agents (2D)
- Inhibitors of the renin-angiotensin system may be more effective in the minimisation of proteinuria but they should be used with caution in the first 3 months post transplant (2C)
- Resistant hypertension may be due to transplant renal artery stenosis and should be investigated according to local practice (2D)

Audit Measures

- The proportion of patients receiving a target blood pressure of 140/90mmHg or 130/80mmHg in the presence of proteinuria PCR>100 or ACR>70)
- Proportion of patients with proteinuria assessed by dipstick and, if present, quantified at each clinic visit

Rationale

The goal of treatment of hypertension in KTR is to reduce the risk of the cardiovascular complications of hypertension (stroke, heart failure, myocardial infarction and arrhythmia) and to preserve or minimise the rate of long term decline in graft function. Blood pressure targets and strategies for treatment should be tailored to the individual patient taking into account proteinuria, the presence of end organ damage such as left ventricular hypertrophy, potential side effect profile of antihypertensive therapy, immunosuppression, and any other relevant lifestyle factors such as safety in pregnancy in female KTRs of childbearing age.

Hypertension is common following kidney transplantation and is associated with reduced graft and patient survival. Many patients require treatment with more than one antihypertensive agent and despite therapy many patients fail to meet treatment targets. Hypertension is a side effect of immunosuppressive therapy, in particular corticosteroids and calcineurin inhibitors, and may also be a consequence of poor graft function leading to salt and water retention. Defining the target optimal target blood pressure in KTRs is challenging and is mainly based on observations made from retrospective registry studies. When office/clinic blood pressure is felt to be unrepresentative of actually blood pressure (such as suspected 'white coat' hypertension, resistant hypertension), home recordings or ambulatory monitoring can be useful, but most studies demonstrate lower measures with these methods and revised targets should be set¹¹⁵. Resistant hypertension, often defined in the general population as clinic BP >160/90 mmHg on 3 or more antihypertensive agents is common in KTR and suggests that further work up is required to exclude transplant renal artery stenosis, non-compliance, or rarer causes of secondary hypertension e.g. primary aldosteronism or pheochromocytoma. In the general population, there has been a move towards more frequent use of ambulatory blood pressure monitoring or home blood pressure measurements for the diagnosis and treatment of hypertension. This is a reasonable strategy for KTRs with the proviso that KTRs are known to be at higher risk of the complications of hypertension and there should be a low threshold for therapeutic intervention when blood pressure targets are exceeded. Where resistant hypertension is suspected, target end organ damage should be assessed using echocardiography to look for the presence of left ventricular hypertrophy.

There are no large scale trials of any single antihypertensive agent in KTRs. Treatment strategies are therefore either defined from extrapolation from other populations or from retrospective analysis of registry data or post hoc studies from clinical trials. The use of blockers of the renin-angiotensin system has been associated with improved patient and graft survival in some but not all retrospective studies, and effectively reduces proteinuria in KTRs although recent trial data suggest that this may not translate into better longer term graft outcomes¹¹⁶⁻¹¹⁸. However, this may be balanced against the risk of hyperkalaemia, more anaemia, and the drop of GFR when these agents are started. The drop in GFR may be predictable and if non progressive and <30% may reflect simply the haemodynamic effect of starting these agents rather than any more serious consequence of these agents. A greater drop in GFR is suggestive of transplant renal artery stenosis and may prompt further imaging. The use of dihydropyridine calcium antagonists may also have benefits in hypertension associated with use of CNIs. Modification of immunosuppression including switching from ciclosporin to tacrolimus, minimisation of calcineurin inhibitors, switching to CNI-free immunosuppression and withdrawal of corticosteroids may all be associated with lower blood pressure. All these strategies may be considered in the presence of resistant hypertension in the absence of any other cause, when graft function is stable and there have been no recent rejection episodes.

References

115. Wadei HM, Textor SC. Hypertension in the kidney transplant recipient. *Transplant Rev (Orlando)*. 2010 Jul;24(3):105-20
116. Knoll GA, Fergusson D, Chasse M, et al. Ramipril versus placebo in kidney transplant patients with proteinuria: a multicentre, double-blind, randomised controlled trial. *The lancet Diabetes & endocrinology* 2015.
117. Opelz G, Dohler B. Cardiovascular death in kidney recipients treated with renin-angiotensin system blockers. *Transplantation* 2014;97:310-5.
118. Tutone VK, Mark PB, Stewart GA, et al. Hypertension, antihypertensive agents and outcomes following renal transplantation. *Clinical transplantation* 2005;19:181-92.

Guideline 6.2 – KTR: Dyslipidaemia

We suggest that the management of dyslipidaemia take into account that:

- Fasting lipid levels should be measured on an annual basis in all renal transplant recipients (2C)
- Treatment targets should be the same as in the general population (2C)
- KTRs at increased primary or secondary CV risk receive statin therapy to reduce the risk of coronary artery disease (2C)
- The choice and dose of statin should take into account concurrent immunosuppression. High dose simvastatin (≥ 40 mg daily) should be avoided in conjunction with ciclosporin and/calcium channel antagonists (2D)

Audit Measures

- Proportion of renal transplant patients with measure of lipids
- Proportion of renal transplant patients taking statins for primary and secondary prevention of cardiovascular disease
- Proportion of transplant patients receiving other lipid lowering treatments
- Proportion of patients achieving dyslipidaemia targets

Rationale

Patients with CKD who have reached end stage renal disease (ESRD) requiring transplantation are at high cardiovascular risk. The leading cause of graft loss is death with a functioning graft, whilst the leading cause of death in renal transplant recipients is cardiovascular disease. Therefore, it is imperative that cardiovascular risk is lowered aggressively to optimise patient and graft outcomes. Kidney transplant recipients have a high prevalence of dyslipidaemia, including raised total, HDL and LDL cholesterol and hypertriglyceridaemia¹¹⁹. Dyslipidaemia is a consequence of immunosuppressive therapy, specifically corticosteroids, ciclosporin (more so than tacrolimus), sirolimus and everolimus¹²⁰. Lipid lowering therapy is likely to be beneficial for many renal transplant recipients. Unlike previous versions of these guidelines, we have taken account of the recent JBS3 guidelines, which suggest measurement of *non-fasting* lipid profile.¹²¹ It seems sensible to use similar clinical practice for assessment of lipid status in KTR as in the general population. However, the JBS3 calculator is not likely to be appropriate for use in KTR and other calculators specific to KTR exist¹²². In keeping with the ethos of lifetime risk rather than an absolute LDL-cholesterol value being used as a trigger for intervention with lipid lowering therapy, we suggest that a 10% 7-year risk of a major adverse cardiac event using the risk calculator derived from the ALERT study, or a 10% 10-year risk of myocardial infarction or stroke using the JBS3 calculator (which is likely to underestimate risk in transplant patients) should be used as a trigger for consideration of lipid lowering therapy.

Performing a baseline assessment of lipid status allows concordance with therapy to be assessed and additionally allows estimation of the level of lipid reduction. Concordance with therapy is recognised to be challenging in renal transplant recipients who are usually on multiple agents often including immunosuppression, antihypertensive therapy and antimicrobial prophylaxis. Lipid assessment should be performed once immunosuppressive drug dosing is stable and the risk of acute rejection requiring corticosteroid therapy has fallen. This is likely to be least three months after transplantation although this will vary with individual patients. Specific targets for treatment have not been recommended as they are difficult to define and are not well defined in the general population. The choice and dose of statin is more likely to be selected on the basis of safety and the immunosuppressive regimen rather than potency in lowering LDL cholesterol.

Statins have similar effects on the secondary dyslipidaemia seen in renal transplant recipients as is demonstrated in primary dyslipidaemia in the general population. The ALERT study showed in a large scale randomised controlled trial that long-term treatment with fluvastatin (40–80 mg per day) compared with placebo non-significantly reduced the risk of coronary death or non-fatal MI in ciclosporin-treated renal transplant recipients^{123,124}. In this study, fluvastatin led to a significant 35% relative reduction in the risk of cardiac death or non-fatal MI. 18.7% patients in ALERT were diabetic at baseline and diabetes was a risk factor for cardiac death in this study¹²⁵. However, there was not a significant reduction in cardiac events in diabetic renal transplant patients.

Statins are metabolised by the cytochrome P450 microsomal enzyme system and concurrent therapy with inhibitors of this system such as ciclosporin or tacrolimus can lead to greater statin exposure and higher risk of side effects such as rhabdomyolysis. This risk appears to be greater with simvastatin and is lowest with fluvastatin or pravastatin^{126,127}. Ezetimibe appears to be safe in renal transplant recipients; although it has been reported to interfere with ciclosporin levels, recent reports suggest this is unlikely to be a major clinical problem^{128,129}. Fibrates have a high risk of side effects and are best avoided in renal transplant recipients.

References

119. Jardine AG, Gaston RS, Fellstrom BC, Holdaas H. Prevention of cardiovascular disease in adult recipients of kidney transplants. *Lancet* 2011;378:1419-27.
120. Miller LW. Cardiovascular toxicities of immunosuppressive agents. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2002;2:807-18.

121. Board JBS. Joint British Societies' consensus recommendations for the prevention of cardiovascular disease (JBS3). *Heart* 2014;100 Suppl 2:ii1-ii67.
122. Soveri I, Holme I, Holdaas H, Budde K, Jardine AG, Fellstrom B. A cardiovascular risk calculator for renal transplant recipients. *Transplantation* 2012;94:57-62.
123. Holdaas H, Fellstrom B, Cole E, et al. Long-term cardiac outcomes in renal transplant recipients receiving fluvastatin: the ALERT extension study. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2005;5:2929-36.
124. Holdaas H, Fellstrom B, Jardine AG, et al. Effect of fluvastatin on cardiac outcomes in renal transplant recipients: a multicentre, randomised, placebo-controlled trial. *Lancet* 2003;361:2024-31.
125. Jardine AG, Fellstrom B, Logan JO, et al. Cardiovascular risk and renal transplantation: post hoc analyses of the Assessment of Lescol in Renal Transplantation (ALERT) Study. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2005;46:529-36.
126. Riella LV, Gabardi S, Chandraker A. Dyslipidemia and its therapeutic challenges in renal transplantation. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2012;12:1975-82.
127. Stirling CM, Isles CG. Rhabdomyolysis due to simvastatin in a transplant patient: Are some statins safer than others? *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2001;16:873-4.
128. Buchanan C, Smith L, Corbett J, Nelson E, Shihab F. A retrospective analysis of ezetimibe treatment in renal transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2006;6:770-4.
129. Turk TR, Voropaeva E, Kohnle M, et al. Ezetimibe treatment in hypercholesterolemic kidney transplant patients is safe and effective and reduces the decline of renal allograft function: a pilot study. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2008;23:369-73.

Guideline 6.3 – KTR: Diabetes mellitus

We suggest that the detection and treatment of diabetes mellitus in KTR should include the following:

- Screening for the development of post transplant diabetes mellitus (PTDM) by dipstick urinalysis and measurement of blood sugar level at each clinic visit (2C)
- Post transplant immunosuppression should take into account risk factors for the development of diabetes. (2C)
- We recommend that diagnosis of PTDM is made based on WHO criteria for diagnosis of diabetes mellitus based on fasting or random blood, serum HBA1c or oral glucose tolerance testing (1C)
- We suggest that a diagnosis of PTDM is made once patients are established on stable maintenance immunosuppression (2D)
- We suggest that post-transplant diabetes should be managed in collaboration with specialists in diabetic medicine (2D)
- All units should have a protocol for the management of post-transplant diabetes (2C)
- KTR with diabetes (either prior to transplantation or PTDM) should undergo screening for diabetic complications (retinal screening, foot care, neuropathy) in line with guidelines for non KTR patients with diabetes (2D)

Audit Measures

- Incidence of post transplant diabetes mellitus (PTDM) at three months and at annual intervals thereafter
- Proportion of patients who require insulin, and in whom remedial action is undertaken – minimisation of steroids and switching of CNIs
- The proportion of patients with PTDM enrolled in screening for extra-renal complications of PTDM

Rationale

Post-transplant diabetes mellitus (PTDM) is common following successful kidney transplantation and occurs in 5-20% of KTRs. PTDM impacts on patient survival, particularly by increasing the risk of cardiovascular disease. Data are limited on the exact magnitude of increase in cardiovascular risk associated with PTDM or the risk of microvascular complications of PTDM (including retinopathy, neuropathy and late graft failure due graft diabetic nephropathy). Risk factors for PTDM are well established and include increasing age, obesity, known glucose intolerance or metabolic syndrome, hepatitis C and family history. Transplant-specific risk factors for PTDM include tacrolimus use (compared to ciclosporin) and corticosteroids. Whilst a number of studies and prior guidelines have advocated switching immunosuppression specifically to minimise the risk of PTDM, this should be balanced against the need for optimal graft function and risk of acute rejection and subsequent need for further steroids, which may paradoxically increase PTDM risk. Similarly, steroid minimisation is an attractive strategy for reducing the risk of PTDM, but published evidence does not consistently support this.

Many patients will exhibit transient hyperglycaemia in the first month after transplantation due to a combination of high dose corticosteroids and CNI inhibition. Whilst this identifies a group of patients at risk of PTDM, it is not useful to over diagnose PTDM and it is reasonable to confirm the presence of PTDM once immunosuppression is stable, typically at 3 months post transplantation.

Once diagnosed, care of the KTR with PTDM should be co-ordinated in a similar fashion to patients with diabetes without transplants. Particular focus should be paid to maintenance of good glycaemic control, treatment of conventional cardiovascular risk factors and screening for the microvascular complications of PTDM. There is no current evidence for any particular hypoglycaemic in KTRs and lifestyle measures, oral hypoglycaemic agents and insulin can all be used to address glycaemic control.

Guideline 6.4 – KTR: Ischaemic heart disease

- We suggest that KTRs receive standard treatment for ischaemic heart disease including thrombolysis, revascularisation, and secondary prevention (2C)

Audit Measures

- Proportion of patients with ischaemic heart disease
- Proportion of patients suffering myocardial infarction
- Proportion of patients undergoing primary revascularisation
- Proportion of patients receiving secondary prevention with a statin, anti-platelet agents and RAS blockers

Rationale

Coronary artery disease (CAD) is common in patients with ESRD, including transplant recipients, and is a major contributory factor to cardiovascular mortality and morbidity. The presence of coronary artery disease often influences the decision to transplant list patients with ESRD. Irrespective of the presence or severity of pre-transplant CAD, once transplanted, CAD is reported in 14–20% (previous myocardial infarction MI or CAD)¹³⁰⁻¹³². Post transplantation myocardial infarction (MI) is common, affecting approximately 11.1% of patients by 3 years post transplantation, and much of this risk is experienced early, within the first 6 months of transplantation¹³³. MI risk has been shown to be linked with modifiable factors including delayed graft function, post transplantation diabetes and graft failure, and in turn, MI predicts graft failure and death. Additionally, the ALERT trial¹²⁴ demonstrated that determinants of non-fatal myocardial infarction in RTR include total cholesterol level, prior CAD and previous acute rejection. Combined, these data suggest that whilst RTR share common risk factors with the general population for CAD and MI post transplantation, there

are further graft-specific aspects to post-transplant CAD. It is known that patients with ESRD, including transplant recipients, are less likely to undergo cardiac intervention (bypass grafting, thrombolytics or per-catheter therapies), possibly because of higher complication rates, and are less likely to receive secondary prevention. There is no reason to believe that transplant recipients will benefit less than patients in the general population, many of whom have renal impairment. Patients with renal transplants should have equal access to cardiac investigations and surgery as patients without CKD.

References

130. Carpenter MA, Weir MR, Adey DB, House AA, Bostom AG, Kusek JW. Inadequacy of cardiovascular risk factor management in chronic kidney transplantation - evidence from the FAVORIT study. *Clinical transplantation* 2012;26:E438-46.
131. Collins AJ, Foley RN, Chavers B, et al. US Renal Data System 2013 Annual Data Report. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2014;63:A7.
132. Pilmore HL, Skeans MA, Snyder JJ, Israni AK, Kasiske BL. Cardiovascular disease medications after renal transplantation: results from the Patient Outcomes in Renal Transplantation study. *Transplantation* 2011;91:542-51.
133. Lentine KL, Brennan DC, Schnitzler MA. Incidence and predictors of myocardial infarction after kidney transplantation. *Journal of the American Society of Nephrology : JASN* 2005;16:496-506.

Guideline 6.5 - KTR: Smoking cessation

We recommend that smoking should be discouraged in transplant recipients (1A)

Audit Measures

- Proportion of KTRs who smoke
- Proportion of cigarette smoking KTRs who have been given formal advice or offer help with cessation

Rationale

Cigarette smoking is strongly associated with the reduced life expectancy, several forms of malignancy, respiratory disease and premature cardiovascular disease in the general population. Whilst the evidence is less comprehensive in KTRs, cigarette smoking has been shown to be associated with reduced patient survival, malignancy, and increased cardiovascular events¹³²⁻¹³⁴. In the general population, various intervention strategies have been shown to be beneficial in encouraging smoking cessation (nicotine replacements - gum, patch, and inhaled, counselling, and Bupropion)^{135,136}. Maintenance of abstinence is challenging, as 10-20% of former smokers will resume smoking following previously successful cessation¹³⁷. The long-term benefits of smoking cessation have not been proven in transplant recipients, nor are long-term studies likely to be performed. However, strategies for smoking cessation are safe and likely to produce the same benefits seen in other populations or public health studies. A local strategy should be available and record made of the advice given and available.

References

134. Lentine KL, Brennan DC, Schnitzler MA. Incidence and predictors of myocardial infarction after kidney transplantation. *Journal of the American Society of Nephrology : JASN* 2005;16:496-506.
135. Danpanich E, Kasiske BL. Risk factors for cancer in renal transplant recipients. *Transplantation* 1999;68:1859-64.
136. Hurst FP, Altieri M, Patel PP, et al. Effect of smoking on kidney transplant outcomes: analysis of the United States Renal Data System. *Transplantation* 2011;92:1101-7.
137. Eisenberg MJ, Filion KB, Yavin D, et al. Pharmacotherapies for smoking cessation: a meta-analysis of randomized controlled trials. *CMAJ* 2008;179:135-44.

138. Strassmann R, Bausch B, Spaar A, Kleijnen J, Braendli O, Puhan MA. Smoking cessation interventions in COPD: a network meta-analysis of randomised trials. *Eur Respir J* 2009;34:634-40.
139. Corbett C, Armstrong MJ, Neuberger J. Tobacco smoking and solid organ transplantation. *Transplantation* 2012;94:979-87.

Guideline 6.6 KTR: Lifestyle measures

We suggest that advice on healthy lifestyle forms a routine part of post-transplant care:

- Maintenance of a healthy diet should be encouraged (2C)
- An ideal weight should be targeted (body mass index (BMI) ≤ 25 kg/m²) (2C)
- Weight management services should be available (2C)
- We suggest that KTRs participate in physical activity at a level similar to that recommended to age and co-morbidity matched counterparts from the general population (2D)
- Alcohol consumption should be within national guidelines (2D)
- Recreational drug use should be avoided (2D)
- The use of over-the-counter medications (without discussion with clinical staff) and non-proprietary medications (e.g. herbal medicines) should be discouraged (2D)

Audit Measures

- Proportion of patients who are obese

Rationale

Transplant recipients have often been subjected to dietary restriction associated with advanced CKD, removal of which after transplantation is one of the factors contributing to weight gain, the metabolic syndrome, diabetes and their sequelae. Whilst metabolic syndrome and obesity have been associated with poorer graft outcomes, the overall impact on obesity on patient and graft outcomes is less than might be expected¹⁴⁰. However, as longer follow up data emerge from the general population regarding the association between obesity and cardiovascular disease and malignancy¹⁴¹ it would seem prudent for KTRs to avoid obesity. KTR should have access to dietary advice, and to weight management services if necessary. Pharmacological intervention for obesity has not been assessed in a clinical trial in KTR and may interfere with the metabolism and absorption of immunosuppressive agents¹⁴². Bariatric surgery is similarly unproven in this population and likely to have a higher incidence of side effects and potential interactions. Dose reduction or withdrawal of corticosteroids helps weight loss but more intensive monitoring is essential around the time of dose changes.

There are very limited data on any specific dietary interventions in KTRs. However, as premature cardiovascular disease is a leading cause of death following transplantation, we suggest that patients follow a diet, which minimises intake of saturated fat, sugar and salt.

We suggest that in addition to body weight control, dietary advice should be given to ensure adequate calcium and magnesium intake and control of serum phosphate level.

We suggest that KTRs avoid dietary intake of produce associated with an increased risk of infections such as *listeria monocytogenes*, *campylobacter jejuni*, etc, e.g. raw shellfish, unpasteurised dairy products. Grapefruit juice should be avoided due to the potential for interference in the metabolism of immunosuppressant drugs (tacrolimus, ciclosporin) by intestinal CYP3A4, leading to increased drug levels.

KTRs have often been restricted in their ability to keep physically active whilst on dialysis. Maintaining a 'healthy' level of physical activity is likely to be beneficial and exercise based interventions have been shown to have a positive impact on quality of life and aerobic capacity in KTRs¹⁴³. Participation in sporting events is often beneficial. Due to the position or the transplant, participation in sports where a direct blow to the allograft is possible is not recommended (e.g. kickboxing).

Alcohol abuse occurs in a small proportion of KTRs though the prevalence and severity of alcohol misuse is difficult to define. Alcohol use within recommended guidelines after transplantation is likely to be safe, whilst alcohol or substance abuse are associated with an increased of premature death^{144,145}. Access to counselling, addiction services and rehabilitation should be available.

Non-steroidal anti-inflammatory drugs are widely available 'over the counter' and should be avoided. There are potential interactions with OTC and "herbal medications" (e.g. St. John's Wort). Patients should be aware of the increased risk and potential sequelae of drug interactions, and encouraged to discuss any proposed changes to medication with clinical staff or an expert renal pharmacist.

References

- 140. Nicoletto BB, Fonseca NK, Manfro RC, Goncalves LF, Leitao CB, Souza GC. Effects of obesity on kidney transplantation outcomes: a systematic review and meta-analysis. *Transplantation* 2014;98:167-76.
- 141. Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet* 2014;384:755-65.
- 142. Evans S, Michael R, Wells H, et al. Drug interaction in a renal transplant patient: cyclosporin-Neoral and orlistat. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2003;41:493-6.
- 143. Heiwe S, Jacobson SH. Exercise Training in Adults With CKD: A Systematic Review and Meta-analysis. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2014;64:383-93.
- 144. Machnicki G, Pinsky B, Takemoto S, et al. Predictive ability of pretransplant comorbidities to predict long-term graft loss and death. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:494-505.
- 145. Parker R, Armstrong MJ, Corbett C, Day EJ, Neuberger JM. Alcohol and substance abuse in solid-organ transplant recipients. *Transplantation* 2013;96:1015-24.

7. Kidney Transplant Recipient (KTR): Neoplasia (Guidelines 7.1-7.3)

General Concepts

Neoplasia is more common in KTRs due to impaired immunosurveillance. In particular, virally driven cancers are more prevalent e.g. human papilloma (HPV)-induced cervical cancer (see Table 3). The relative risk of cancer is higher in younger patients (20x relative risk of neoplasia) than older patients (2x for over 65s)^{144,145}. KTRs with neoplasia have worse outcomes than members of the general population, probably due to increased toxicity from treatment. Preventative strategies are therefore paramount, which may involve screening and the minimisation or modification of immunosuppressive therapy. If cancer develops then part of the treatment will involve reducing and/or modifying immunosuppressive therapy. This is likely to be more beneficial in those cancers with higher relative risks in KTRs (e.g. more likely to have a clinical impact in non-melanoma skin cancer than pancreatic cancer). Emerging evidence supports the notion that low dose immunosuppression and the use of m-TORi may reduce the incidence and recurrence of some cancers¹⁴⁶⁻¹⁴⁸.

References

- 146. NHS Cancer Screening Programmes. 2014. at <http://www.cancerscreening.nhs.uk/index.html>.)
- 147. Wong G, Howard K, Webster A, Chapman JR, Craig JC. The health and economic impact of cervical cancer screening and human papillomavirus vaccination in kidney transplant recipients. *Transplantation* 2009;87:1078-91.

148. Cherikh WS, Kauffman HM, McBride MA, Maghirang J, Swinnen LJ, Hanto DW. Association of the type of induction immunosuppression with posttransplant lymphoproliferative disorder, graft survival, and patient survival after primary kidney transplantation. *Transplantation* 2003;76:1289-93.

149. Kauffman HM, Cherikh WS, Cheng Y, Hanto DW, Kahan BD. Maintenance immunosuppression with target-of-rapamycin inhibitors is associated with a reduced incidence of de novo malignancies. *Transplantation* 2005;80:883-9.

150. Salgo R, Gossmann J, Schofer H, et al. Switch to a sirolimus-based immunosuppression in long-term renal transplant recipients: reduced rate of (pre-)malignancies and nonmelanoma skin cancer in a prospective, randomized, assessor-blinded, controlled clinical trial. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2010;10:1385-93.

Relative risk	KTR Common cancers
High RR >5	Kaposi's sarcoma Eye Lymphoma Kidney Non-melanoma skin Lip Thyroid
Medium RR 1-5	Melanoma Cervix Vulvovaginal Bladder Colon Lung Stomach Oesophagus Oropharynx and Larynx Myeloma Anus Leukaemia Hepatobiliary
No increase	Breast Prostate Ovary Uterus Pancreas Brain Testis

Table 3- Association of risk of common malignancies in KTRs

Guideline 7.1 – KTR: Screening for cancer

We suggest that the organisation of screening for neoplasia in KTRs take into account:

- Screening should be similar to the general population for cervical, breast, colon and prostate cancer (2C)
- Screening is not recommended for renal cell carcinoma (2C)
- Patient education pre and post transplantation (1C)
- Patients should be aware of malignancy risk and encouraged to report symptoms which may represent *de novo* malignancy (e.g. breast or testicular lumps) (2D)

- Skin surveillance by a healthcare professional at least biannually up to 5 years post-transplant and annually from 5 years post-transplant (2C)
- Patients with cirrhosis should undergo an annual hepatic ultrasound and determination of serum alpha feto-protein (2C)

Audit Measure

- Proportion of patients having screening procedures for neoplasia at the annual review clinic

Rationale

The merits of any screening programme must balance the individual's risk of developing the disease, their prognosis if detected, and the risk of harm from screening. Screening should be individualised and reflect co-morbidities and other competing risks (e.g. vascular disease). Some authors have advocated more frequent screening (e.g. annual cervical screening) but there is little evidence to support these assertions. Thus screening should follow the pattern in the general population for most common cancer¹⁴⁵. The national cancer screening protocols are described on the NHS Cancer Screening Programme (NHSCSP) website¹⁴⁶. Patient education regarding neoplasia should be undertaken pre-transplant, with reinforcement post-transplant. Education should include breast, testicular and skin self-examination. Information about risk factors for the development of NSMC should also be included¹⁵¹.

That all KTRs should undergo skin surveillance is not in doubt but the frequency of survey to optimally balance early detection and treatment with finite resource is unclear¹⁵². The incidence of non-melanoma skin cancers increases with duration post-transplant and skin surveillance should therefore be more frequent^{151,153}. Encouragement of self-examination and education should be reinforced in the initial years following transplant, supplemented with skin surveillance by a trained healthcare professional at least biannually¹⁵⁴. From 5 years post-transplant, skin surveillance should be undertaken annually to reflect the increased likelihood of developing NSMC¹⁵³. Smoking cessation should be encouraged in all KTRs. A formal protocol for the management of smoking cessation should be available in each transplant centre (see Guidelines 6.5)

References

151. Harden PN, Fryer AA, Reece S, Smith AG, Ramsay HM. Annual incidence and predicted risk of nonmelanoma skin cancer in renal transplant recipients. *Transplantation proceedings* 2001;33:1302-4.
152. Garg S, Carroll RP, Walker RG, Ramsay HM, Harden PN. Skin cancer surveillance in renal transplant recipients: re-evaluation of U.K. practice and comparison with Australian experience. *Br J Dermatol* 2009;160:177-9.
153. Ramsay HM, Reece SM, Fryer AA, Smith AG, Harden PN. Seven-year prospective study of nonmelanoma skin cancer incidence in U.K. renal transplant recipients. *Transplantation* 2007;84:437-9.
154. Stasko T, Brown MD, Carucci JA, et al. Guidelines for the management of squamous cell carcinoma in organ transplant recipients. *Dermatol Surg* 2004;30:642-50.

Guideline 7.2 – KTR: Non-Melanoma Skin Cancer (NMSC)

We recommend that KTRs should be educated about the adverse effects of solar exposure (1C)

Guideline 7.3 – KTR: Non-Melanoma Skin Cancer (NMSC)

We suggest that KTRs that an individualised assessment of hazard should be made according to risk factors (2C)

Guideline 7.4 – KTR: Non-Melanoma Skin Cancer (NMSC)

We recommend that patients should be encouraged to cover their skin in direct sunlight and to use total sunblock (Sun Protection Factor \geq 50) (1D)

Guideline 7.5 – KTR: Non-Melanoma Skin Cancer (NMSC)

We suggest that self-examination should be encouraged with guidance provided. This should be supplemented by at least biannual review by a trained healthcare professional up to 5 years post-transplant and annual review from 5 years (2C)

Guideline 7.6 – KTR: Non-Melanoma Skin Cancer (NMSC)

We suggest that the prescription of acitretin as chemoprophylaxis be considered in those with \geq 2 previous NMSC if there are no contraindications (2B)

Rationale

Certain patient groups are at higher risk of non-melanoma skin cancer, particularly the fair-skinned living in a sunny climate. Other risk factors include occupation, behaviour, previous skin cancer, childhood solar exposure and family history. It is sensible to minimise exposure and use sun block. Acitretin (0.2-0.4 mg/kg/day) may reduce total NSMCs in those who have had \geq 2 previous NSMCs and thus use should be considered in those with previous skin cancer¹⁶¹. There is some evidence that sirolimus reduces the incidence of second tumours but at the expense of increased side effects and possibly adverse effects on graft function^{155,156,162}. Recent studies suggest that m-TORi may be associated with fewer NMSC, particularly cutaneous Kaposi sarcoma and recent data suggest that switching KTRs to sirolimus is associated with reduction of secondary NMSC^{156,163}. The role of HPV vaccination in KTRs is unclear, but it is an inactivated vaccine that can be administered safely either before or after transplantation¹⁶⁴.

References

155. Kauffman HM, Cherikh WS, Cheng Y, Hanto DW, Kahan BD. Maintenance immunosuppression with target-of-rapamycin inhibitors is associated with a reduced incidence of de novo malignancies. *Transplantation* 2005;80:883-9.
156. Salgo R, Gossmann J, Schofer H, et al. Switch to a sirolimus-based immunosuppression in long-term renal transplant recipients: reduced rate of (pre-)malignancies and nonmelanoma skin cancer in a prospective, randomized, assessor-blinded, controlled clinical trial. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2010;10:1385-93.
157. Harden PN, Fryer AA, Reece S, Smith AG, Ramsay HM. Annual incidence and predicted risk of nonmelanoma skin cancer in renal transplant recipients. *Transplantation proceedings* 2001;33:1302-4.
158. Garg S, Carroll RP, Walker RG, Ramsay HM, Harden PN. Skin cancer surveillance in renal transplant recipients: re-evaluation of U.K. practice and comparison with Australian experience. *Br J Dermatol* 2009;160:177-9.
159. Ramsay HM, Reece SM, Fryer AA, Smith AG, Harden PN. Seven-year prospective study of nonmelanoma skin cancer incidence in U.K. renal transplant recipients. *Transplantation* 2007;84:437-9.
160. Stasko T, Brown MD, Carucci JA, et al. Guidelines for the management of squamous cell carcinoma in organ transplant recipients. *Dermatol Surg* 2004;30:642-50.
161. Kadakia KC, Barton DL, Loprinzi CL, et al. Randomized controlled trial of acitretin versus placebo in patients at high-risk for basal cell or squamous cell carcinoma of the skin (North Central Cancer Treatment Group Study 969251). *Cancer* 2012;118:2128-37.
162. Stallone G, Schena A, Infante B, et al. Sirolimus for Kaposi's sarcoma in renal-transplant recipients. *The New England journal of medicine* 2005;352:1317-23.
163. Euvrard S, Morelon E, Rostaing L, et al. Sirolimus and secondary skin-cancer prevention in kidney transplantation. *The New England journal of medicine* 2012;367:329-39

164 Kumar D, Unger ER, Panicker G, Medvedev P, Wilson L, Humar A. Immunogenicity of quadrivalent human papillomavirus vaccine in organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2013;13:2411-7.

Guideline 7.7 – KTR: Immunosuppression in cancers

We suggest that the overall level of immunosuppression should be reduced if neoplasia develops (2C)

Guideline 7.8 – KTR: Immunosuppression in cancers

We suggest that m-TORis are considered as alternative immunosuppressive agents in KTRs who develop *de novo* malignancy (2C)

Guideline 7.9 – KTR: Immunosuppression in Kaposi’s sarcoma

We suggest that m-TORis has specific anti-tumour effects in Kaposi’s sarcoma and switching to this medication should be considered (2C)

Rationale

There is no evidence that any particular immunosuppressant agent is linked to a particular cancer other than the association of TDAs and PTLD^{28,146}. It is generally agreed that the level of immunosuppression should be decreased when cancer occurs. This decision should be individualised according to the stage of the cancer at diagnosis, the likely impact of a reduction in immunosuppression, the availability of treatment for the tumour, and potential drug interactions between the chemotherapy and immunosuppressive agents. In general, the effect of reducing the overall level of immunosuppression is more likely to be beneficial where the relative risk of the tumour in KTRs is higher. m-TORis are indicated in the treatment of Kaposi sarcoma in addition to reduced levels of overall immunosuppression¹⁵⁴.

8. Kidney Transplant Recipient (KTR): Infection Complications (Guidelines 8.1-8.7)

Guideline 8.1 – KTR: Vaccination

Guideline 8.1.1 – KTR: Vaccination

We recommend that KTRs:

- Should be vaccinated with inactivated viruses as per the normal population (1D)
- Should receive annual influenza vaccination unless contraindicated (1C)

Guideline 8.1.2 – KTR: Vaccination

We suggest that KTRs:

- Should have HBsAb levels rechecked annually and consider revaccination if antibody titres fall below 10mIU/ml (2D)
- Should not receive live attenuated vaccines (2C)
- Should receive pneumococcal vaccine and a booster every five years (2D)

Rationale

Vaccination for KTRs and their household members should ideally be completed prior to transplantation. A minimum of 4 weeks is recommended between vaccination with live attenuated vaccines and transplantation. After transplantation, there is no evidence to link vaccination with rejection. After transplantation it is safe to administer inactivated vaccines but live attenuated vaccines should be avoided (see Table 4 below) as small studies have demonstrated concerns regarding their safety and efficacy in KTRs. Vaccination should probably be carried out at least 3 months and preferably 6 months after transplantation when the immunosuppression has been reduced. Consideration should be given to vaccination of close household contacts where appropriate e.g. Varicella vaccination for children of VZV seronegative KTRs.

Inactive Vaccine	Live Attenuated Vaccine
Inactivated Influenza	Varicella Zoster
Hepatitis A	Mumps
Hepatitis B	Rubella
Inactivated Polio	Measles
Diphtheria	BCG
Tetanus	Smallpox
Meningococcal	Yellow Fever
Pneumococcal	Oral Salmonella
Human Papilloma Virus	Oral Polio
Rabies	
Anthrax	
Intramuscular Salmonella	
Japanese encephalitis	
Inactivated intravenous cholera vaccine	

Table 4. Commonly used inactive and live attenuated vaccines

The HPV vaccine has been assessed in a small study of transplant recipients. Although the vaccine was safe and tolerated, immunogenicity was suboptimal¹⁶⁵. There is a strong link between HPV and anogenital and non-melanoma skin cancer and, given the tolerability of these vaccines, some authors have recommended vaccination for all female KTRs aged between 9 and 26¹⁶⁶. Malaria prophylaxis should consist of chloroquine in sensitive areas, but it may increase levels of ciclosporin. Prophylaxis should therefore start two weeks prior to departure to permit monitoring of drug levels. In areas of chloroquine-resistance three options can be used: atovaquone and proguanil; mefloquine; or doxycycline. The choice of agent will be dictated by local preference and side effect profile but drugs should be started a few weeks prior to departure to allow for checks of renal and hepatic function, full blood count and immunosuppression levels. More extensive guidance is available for KTRs who are travelling overseas¹⁶⁷.

References

165. Kumar D, Unger ER, Panicker G, Medvedev P, Wilson L, Humar A. Immunogenicity of quadrivalent human papillomavirus vaccine in organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2013;13:2411-7.
166. Chow J, Golan Y. Vaccination of solid-organ transplantation candidates. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2009;49:1550-6.
167. Health Information for overseas travel - Department of Health. 2015.

Guideline 8.2 – KTR: Cytomegalovirus disease

Guideline 8.2.1 – KTR: prophylaxis and treatment of CMV disease

We recommend:

- Prophylaxis should be continued for 3-6 months, until immunosuppression has been reduced to long-term maintenance level (1B)
- Treatment should be administered for 6 weeks after treatment with a TDA (1C)

Guideline 8.2.2 – KTR: prophylaxis and treatment of CMV disease

We suggest:

- All transplant units should have the ability to measure CMV serological status and the detection and quantification of viral load (2D)
- Donor and recipient CMV sero-positivity should be recorded at the time of transplantation (2D)
- A written protocolised strategy based either on prophylaxis, or pre-emptive therapy, or both should be implemented (2D)
- For the treatment of mild and moderate CMV disease, oral valganciclovir and intravenous ganciclovir are of equivalent efficacy (2C)
- Treatment of life-threatening CMV disease should be initiated with intravenous ganciclovir (2D)
- Treatment duration should be determined by monitoring viral load (2C)

Audit Measure

- Incidence of CMV disease.

Rationale

CMV infection is the most common serious viral infection affecting renal transplant recipients^{168,169}. It occurs most commonly in CMV naïve recipients of a kidney from a CMV positive donor. However, seropositive transplant recipients may be affected by reactivation of CMV infection and by primary infection with a new genotype. CMV infection is associated with more intensive immunosuppression, treatment of acute rejection episodes and the use of TDAs. For CMV disease there is clear evidence that prophylaxis reduces the severity, delays the onset and prevents CMV infection in CMV negative recipients of CMV positive kidneys^{170,171}. CMV infection is also associated with concomitant infection by other herpes viruses, the significance of which is uncertain but prevention of which may be an added benefit of prophylaxis over pre-emptive strategies¹⁷². CMV prophylaxis is therefore recommended in CMV negative KTRs from a CMV positive donor (D+/R-), and for seropositive KTRs (D-/R+ or D+/R+) exposed to more intensive immunosuppression, in particular TDAs.

In CMV negative recipients of a CMV positive kidney the use of oral antiviral therapy – specifically valaciclovir, aciclovir, ganciclovir, or valganciclovir – is proven to delay the onset of CMV disease, to prevent CMV disease in a proportion of patients, and to limit the severity of disease. It is important to recognise that a proportion of patients will develop CMV disease after stopping prophylaxis and will require clinical and virological monitoring for at least three months following the stopping of prophylactic therapy¹⁷². At present, evidence exists for the use of valaciclovir and valganciclovir^{173,174}. In clinical practice, valaciclovir is not widely available or used, and valganciclovir is the agent of choice. The dose of valganciclovir should be adjusted according to renal transplant function. Most commonly, and based on the available evidence, antiviral prophylaxis is usually continued for 90-180 days following transplantation (depending on D and R status). The rationale is that reduction of immunosuppressive therapy over this period will allow the immune system to combat viral replication once prophylactic therapy is withdrawn^{175,176}.

Treatment of CMV disease is equally effective with either oral valganciclovir or intravenous ganciclovir^{167,168}. However, this study excluded patients with life-threatening CMV infection and it is probably advisable to initiate treatment with intravenous therapy in such circumstances¹⁶⁹.

References

168. Humar A, Snyderman D, Practice ASTIDCo. Cytomegalovirus in solid organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9 Suppl 4:S78-86.
169. Kotton CN, Kumar D, Caliendo AM, et al. Updated international consensus guidelines on the management of cytomegalovirus in solid-organ transplantation. *Transplantation* 2013;96:333-60.
170. Hodson EM, Jones CA, Webster AC, et al. Antiviral medications to prevent cytomegalovirus disease and early death in recipients of solid-organ transplants: a systematic review of randomised controlled trials. *Lancet* 2005;365:2105-15.
171. Lowance D, Neumayer HH, Legendre CM, et al. Valacyclovir for the prevention of cytomegalovirus disease after renal transplantation. *International Valacyclovir Cytomegalovirus Prophylaxis Transplantation Study Group. The New England journal of medicine* 1999;340:1462-70.
172. Humar A, Asberg A, Kumar D, et al. An assessment of herpesvirus co-infections in patients with CMV disease: correlation with clinical and virologic outcomes. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:374-81.
173. van der Beek MT, Berger SP, Vossen AC, et al. Preemptive versus sequential prophylactic-preemptive treatment regimens for cytomegalovirus in renal transplantation: comparison of treatment failure and antiviral resistance. *Transplantation* 2010;89:320-6.
174. Humar A, Lebranchu Y, Vincenti F, et al. The efficacy and safety of 200 days valganciclovir cytomegalovirus prophylaxis in high-risk kidney transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2010;10:1228-37.
175. Luan FL, Stuckey LJ, Park JM, Kaul D, Cibrik D, Ojo A. Six-month prophylaxis is cost effective in transplant patients at high risk for cytomegalovirus infection. *Journal of the American Society of Nephrology : JASN* 2009;20:2449-58.
176. Asberg A, Humar A, Jardine AG, et al. Long-term outcomes of CMV disease treatment with valganciclovir versus IV ganciclovir in solid organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9:1205-13.
177. Asberg A, Humar A, Rollag H, et al. Oral valganciclovir is noninferior to intravenous ganciclovir for the treatment of cytomegalovirus disease in solid organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2106-13.
178. Avery RK. Valganciclovir versus IV ganciclovir for therapy of cytomegalovirus viremia: has victory been achieved? *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2062-3.

Guideline 8.3 – KTR: Epstein Barr virus infection

Guideline 8.3.1 – KTR: EBV infection

We recommend that immunosuppression should be reduced or stopped following the development of PTLD (1C)

Guideline 8.3.2 – KTR: EBV infection

We suggest:

- Both donor and recipient should have their EBV serology recorded at the time of transplantation (2D)
- All high risk (D+/R-) patients (including adults) should have EBV viral load measured immediately after transplantation, monthly for six months, and three monthly to one year (2C)
- EBV viral load should be monitored after the treatment of rejection (2C)
- Total immunosuppression should be reduced when EBV titres rise significantly (2C)

Audit Measures

- Rates of EBV infection and PTLD amongst KTRs
- Completeness of records for EBV donor and recipient serology

Rationale

After transplantation, primary EBV infection may present with a broad spectrum of disorders ranging from asymptomatic infection to high grade non-Hodgkin lymphoma (PTLD). EBV seronegative KTRs, undergoing seroconversion post transplant, are up to 50 times more likely to develop PTLD compared to their seropositive counterparts. The EBV genome is found in more than 90% of PTLD occurring during the first year after transplantation¹⁷⁹. Vigilance is therefore essential, especially since EBV viraemia usually precedes the development of PTLD by 4-16 weeks¹⁸⁰. However, assays for EBV viral load can often be positive in asymptomatic patients (false positives) and so clinical correlation and attention to changes in viral load are essential. Risk factors for early PTLD include primary EBV infection, young donor age, CMV infection, and induction with TDAs. The use of antiviral agents (e.g. valaciclovir or valganciclovir) or immunoglobulins in response to rising viral loads is unproven and cannot be recommended. Since the immune response to EBV infected tissue is thought to depend on EBV-specific T cell responses it is logical to reduce immunosuppressive treatment in the face of clinical EBV infections and PTLD.

References

179. Allen U, Preiksaitis J, Practice ASTIDCo. Epstein-barr virus and posttransplant lymphoproliferative disorder in solid organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9 Suppl 4:S87-96.
180. Rowe DT, Webber S, Schauer EM, Reyes J, Green M. Epstein-Barr virus load monitoring: its role in the prevention and management of post-transplant lymphoproliferative disease. *Transplant infectious disease : an official journal of the Transplantation Society* 2001;3:79-87.

Guideline 8.4 – KTR: Varicella Zoster Virus infection

Guideline 8.4.1 – KTR: VZV infection

We recommend:

- Primary infection (chickenpox) should be treated with intravenous aciclovir or oral valaciclovir until the lesions scab over (1C)
- Uncomplicated shingles should be treated with oral acyclovir or valaciclovir until the lesions scab over (1D)
- Disseminated (>2 dermatomes), ocular or invasive shingles should be treated with intravenous aciclovir until the lesions scab over, together with a reduction in immunosuppression (1B)

- Varicella-susceptible KTRs (i.e. VZV IgG negative) with primary exposure to VZV should receive intravenous immunoglobulins, ideally within 96 hours, but up to a maximum of 10 days following exposure. If unavailable or after 10 days, oral aciclovir should be administered for seven days (1D)

Guideline 8.4.2 – KTR: VZV infection

We suggest:

- Patients on the waiting list who are VZV IgG seronegative should be vaccinated prior to transplantation (2D)
- Immunosuppression should be reduced during primary infection (2D)

Audit Measures

- Annual rates of primary VZV and shingles infection
- Completeness of records for VZV recipient serology in KTRs

Rationale

Acquired by 90% of the population before adulthood, primary infection with VZV causes chickenpox. Thereafter the virus remains latent in the cranial nerve and dorsal root ganglia. Secondary reactivation results in shingles and typical dermatomal blistering skin lesions. Primary infection can be acquired by direct skin contact and by airborne droplet transmission^{181,182}. Primary disease in KTRs can be devastating with severe skin lesions, widespread visceral involvement and disseminated intravascular coagulation. In immune naïve and immunosuppressed individuals, treatment with pooled varicella zoster immunoglobulin has been shown to prevent or ameliorate VZV infection. It seems sensible to vaccinate VZV naïve patients on the waiting list since the vaccine has been shown to be safe¹⁸². As a live attenuated vaccine, VZV vaccine should not be administered post transplant.

References

181. Heining U, Seward JF. Varicella. *Lancet* 2006;368:1365-76.
182. Pergam SA, Limaye AP, Practice ASTIDCo. Varicella zoster virus (VZV) in solid organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9 Suppl 4:S108-15.

Guideline 8.5 – KTR: Herpes Simplex Virus infection

Guideline 8.5.1 – KTR: HSV infection

We recommend:

- Superficial HSV (1 or 2) infection should be treated with appropriate oral agents until the lesions have resolved (1D)
- Systemic HSV infections should be treated with intravenous aciclovir and a reduction in immunosuppression until a response occurs and oral medication continued for at least 14 days (1C)

Guideline 8.5.2 – KTR: HSV infection

We suggest that KTRs suffering frequent recurrent HSV infection should consider oral prophylaxis (2D)

Audit Measure

- Rates and outcomes of HSV infections

Rationale

There is an increased potential for superficial HSV infections to become disseminated or invasive in KTRs. Reactivation most commonly occurs in the first few weeks after transplantation and complicated disease can become life threatening involving multiple organs including the liver, lung and central nervous system. Since treatment is safe and effective, it seems sensible to treat early infections¹⁸³. Due to their gravity, complicated infections should be treated with intravenous therapy and reduction in immunosuppression.

References

183. Razonable RR, Zerr DM, Practice ASTIDCo. HHV-6, HHV-7 and HHV-8 in solid organ transplant recipients. American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons 2009;9 Suppl 4:S97-100.

Guideline 8.6 – KTR: BK nephropathy

Guideline 8.6.1 – KTR: BK nephropathy

We recommend that confirmed BK nephropathy should be treated by reduction in immunosuppression (1D)

Guideline 8.6.2 – KTR: BK nephropathy

We suggest:

- Screening should also be carried out when renal function deteriorates in an unexplained fashion (2D)
- KTRs should be screened for BKV viral load or by performing urine microscopy for decoy cells or by PCR on urine or serum (2C)
- Suspected BK nephropathy should be confirmed by renal biopsy, which should be stained for SV40. Two cores containing medullary tissue should ideally be examined (2D)
- Immunosuppression should be reduced when the serum BKV load exceeds 10^4 copies/ml (2C)
- There is no established specific treatment for BK nephropathy (2D)
- Re-transplantation can safely be considered in patients who have BK nephropathy diagnosed in an earlier graft (2C)

Audit Measures

- Rates of BK viral infection in screening tests
- Rates and outcomes of BK nephropathy

Rationale

The human polyoma BK virus is linked to two major clinical syndromes in KTRs, namely BK nephropathy (BKN) and transplant ureteric stenosis¹⁸⁴⁻¹⁸⁶. BKN occurs in up to 10% of KTRs and is responsible for a significant number (15-50%) of allograft losses. 90% of young adults have serological evidence of prior infection and the DNA virus remains latent in the uroepithelium. The virus becomes active and replicates under the influence of immunosuppression. BK virus is cytolytic and so epithelial cells are shed in the urine as decoy cells and free virus can be detected in the urine. With increased viral replication, BKV spills into the blood and can be detected as BK viraemia by PCR. Approximately half of patients with high level viruria ($>10^7$ copies/mL) will

develop significant BK viraemia (10^4 copies/mL) and half of these will develop histological BKN. Risk factors for BKV include not only both donor and recipient characteristics but also high immunosuppressive burden and intensification of immunosuppression. Definitive diagnosis requires demonstration of the virus in renal tissue, usually by staining with the antibody for large T antigen of SV40. Since the infection can be focal and preferentially affects the renal medulla, two cores including medulla should be examined^{187,188}. The mainstay of treatment is reduction of immunosuppression but there is no evidence that reducing any particular immunosuppressive agent is particularly beneficial¹⁸⁹⁻¹⁹⁰. Common approaches include stopping anti-proliferative agents or reducing CNI levels in the face of rising viral replication¹⁹¹. Specific agents such as intravenous immunoglobulin, quinolones, cidofovir and leflunomide have been shown to have anti-viral activity but there is no definitive evidence to show that they offer any advantage over simply reducing the total immunosuppressive burden¹⁹²⁻¹⁹⁴. Prospective trials are urgently required to explore this question. If a first graft is lost due to BKN there is no evidence that this will adversely affect the outcome of subsequent grafts and no special precautions are necessary (e.g. allograft nephrectomy) prior to re-listing¹⁹⁵.

References

184. Bohl DL, Brennan DC. BK virus nephropathy and kidney transplantation. *Clinical journal of the American Society of Nephrology* : CJASN 2007;2 Suppl 1:S36-46.
185. Dall A, Hariharan S. BK virus nephritis after renal transplantation. *Clinical journal of the American Society of Nephrology* : CJASN 2008;3 Suppl 2:S68-75.
186. Ramos E, Drachenberg CB, Wali R, Hirsch HH. The decade of polyomavirus BK-associated nephropathy: state of affairs. *Transplantation* 2009;87:621-30.
187. Drachenberg CB, Papadimitriou JC, Hirsch HH, et al. Histological patterns of polyomavirus nephropathy: correlation with graft outcome and viral load. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2004;4:2082-92.
188. Hirsch HH, Randhawa P, Practice ASTIDCo. BK virus in solid organ transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2009;9 Suppl 4:S136-46.
189. Hirsch HH, Knowles W, Dickenmann M, et al. Prospective study of polyomavirus type BK replication and nephropathy in renal-transplant recipients. *The New England journal of medicine* 2002;347:488-96.
190. Hardinger KL, Koch MJ, Bohl DJ, Storch GA, Brennan DC. BK-virus and the impact of pre-emptive immunosuppression reduction: 5-year results. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2010;10:407-15.
191. Johnston O, Jaswal D, Gill JS, Doucette S, Fergusson DA, Knoll GA. Treatment of polyomavirus infection in kidney transplant recipients: a systematic review. *Transplantation* 2010;89:1057-70.
192. Maggiore U, Medici MC, Vaglio A, Buzio C. Increased viral load after intravenous immunoglobulin therapy for BK virus-associated nephropathy. *Transplant infectious disease : an official journal of the Transplantation Society* 2010;12:470-2.
193. Pallet N, Burgard M, Quamouss O, et al. Cidofovir may be deleterious in BK virus-associated nephropathy. *Transplantation* 2010;89:1542-4.
194. Randhawa PS, Schonder K, Shapiro R, Farasati N, Huang Y. Polyomavirus BK neutralizing activity in human immunoglobulin preparations. *Transplantation* 2010;89:1462-5.
195. Dharnidharka VR, Cherikh WS, Neff R, Cheng Y, Abbott KC. Retransplantation after BK virus nephropathy in prior kidney transplant: an OPTN database analysis. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2010;10:1312-5.

Guideline 8.7 – KTR: Pneumocystis jirovecii infection- treatment and prophylaxis

We suggest:

- All patients with confirmation (microscopy or PCR) of *Pneumocystis jirovecii* in respiratory secretions should be treated for 14 to 21 days with co-trimoxazole orally or intravenously (15-20mg/kg in three or four divided doses) (2B)
- Patients with contraindications to treatment with co-trimoxazole should receive pentamidine (4mg/kg/day intravenously) (2B)
- Adjunctive glucocorticoid therapy may be considered in patients with severe disease (2D)
- All patients should receive 3-6 months of treatment with co-trimoxazole 480mg daily for *Pneumocystis jirovecii* prophylaxis following renal transplantation (1B)

Rationale

Diagnosis of *Pneumocystis pneumonia* (PCP) requires confirmation by microscopy or PCR from respiratory secretions (induced sputum after inhaled saline or bronchoalveolar lavage). In non-HIV infected individuals, co-trimoxazole has been shown to be effective. Co-trimoxazole has excellent bioavailability and enteral administration is preferred, although intravenous therapy may be necessary in acutely unwell patients. In patients with contraindications to co-trimoxazole, intravenous or aerosolised pentamidine is a suitable alternative although it is associated with a higher rate of side effects. Alternative agents against pneumocystis jirovecii include dapsone or atovaquone. The duration of therapy should be between 14 and 21 days. The use of additional glucocorticoid therapy is not associated with improved survival in non-HIV patients, although treatment with prednisolone has been associated with quicker recovery from severe PCP infection in one small observational study¹⁹⁶. As a result, PCP prophylaxis with co-trimoxazole is recommended and has shown to be beneficial in meta-analysis of 12 randomised studies in 1245 immunocompromised individuals¹⁹⁷.

References

196. Pareja JG, Garland R, Koziel H. Use of adjunctive corticosteroids in severe adult non-HIV *Pneumocystis carinii* pneumonia. *Chest* 1998;113:1215-24.
197. Green H, Paul M, Vidal L, Leibovici L. Prophylaxis of *Pneumocystis pneumonia* in immunocompromised non-HIV-infected patients: systematic review and meta-analysis of randomized controlled trials. *Mayo Clinic proceedings* 2007;82:1052-9.

Guideline 8.8 – KTR: Post-transplant infection prophylaxis

- All patients should receive 3-6 months of treatment with co-trimoxazole 480mg daily (1B)
- Oral antifungal prophylaxis should be administered for one month after transplantation (2C)
- In selected patients, prophylaxis against mycobacterium tuberculosis with daily isoniazid (supplemented with pyridoxine) should be instituted for six months after transplantation (2C)

Rationale

In addition to benefit for PCP, there is good evidence that co-trimoxazole provides effective prophylaxis against urinary tract infections after renal transplantation^{198,199}. Alternatives include cephalosporins and fluoroquinolones. Alternative agents against pneumocystis jirovecii include dapsone, atovaquone or aerosolized pentamidine.

Candida infection is common after renal transplantation and can cause considerable morbidity. It is usually acquired from colonisation of the oral mucosa and so topical oral preparations offer a simple form of prevention without the potential toxicity of systemic preparations.

Tuberculosis in KTRs is usually due to reactivation of quiescent disease under the influence of immunosuppression. In other immunosuppressed populations treatment of latent tuberculosis prevents progression to clinically active tuberculosis. A recent Cochrane Database review recommended prophylaxis with isoniazid (for up to 1 year) in selected recipients at high risk of tuberculosis (based on prior infection or exposure, or patients whose country of origin has a high incidence of tuberculosis infection) with careful monitoring of liver function, especially in patients who are hepatitis B and C positive^{198,200}.

References

198. Currie AC, Knight SR, Morris PJ. Tuberculosis in renal transplant recipients: the evidence for prophylaxis. *Transplantation* 2010;90:695-704.
199. Fox BC, Sollinger HW, Belzer FO, Maki DG. A prospective, randomized, double-blind study of trimethoprim-sulfamethoxazole for prophylaxis of infection in renal transplantation: clinical efficacy, absorption of trimethoprim-sulfamethoxazole, effects on the microflora, and the cost-benefit of prophylaxis. *The American journal of medicine* 1990;89:255-74.
2001. Adamu B, Abdu A, Abba AA, Borodo MM, Tleyjeh IM. Antibiotic prophylaxis for preventing post solid organ transplant tuberculosis. *The Cochrane database of systematic reviews* 2014;3:CD008597.

Guideline 8.9 – KTR: Hepatitis E Virus

We recommend that Hepatitis E Virus (HEV)-screened blood components should be given to all KTR (1C)

Rationale

There has been an increase in the number of reports of cases of hepatitis E virus (HEV) in the United Kingdom. The infection is usually mild and self-limiting in the immunocompetent but there is increasing evidence that HEV infection in patients receiving immunosuppression may lead to persistent infection, which may lead to chronic hepatitis and cirrhosis. HEV may be transmitted through the use of blood and blood products, through transplantation and through diet (especially inadequately cooked pork and pork products such as sausages and offal). Liver tests may be normal or show mild hepatitis. The favoured diagnostic test is by HEV polymerase chain reaction. Liver biopsy may show histology. Conventional management of HEV infection is review immunosuppression and reduce or minimise immunosuppressive burden where possible. If there is no serological evidence of clearance within three months, consider a three-month course of Ribavirin (note this use is off licence)^{201,202}.

References

201. SaBTO/BSBMT. Reducing the risk of Transfusion-transmitted Hepatitis E Virus (HEV) infections in patients undergoing Solid Organ Transplantation (SOT) and Haematopoietic Stem Cell Transplantation (HSCT). 2016.
202. SaBTO/BSBMT. SaBTO Letter to Clinicians - Re Hep E Virus. 2016.

9. Kidney Transplant Recipient (KTR): Bone and Joint disease (Guidelines 9.1-8.7)

Guideline 9.1 KTR: Osteoporosis

We suggest:

- KTRs suffering from osteoporosis or at high potential risk should be considered for steroid-avoiding immunosuppression (2D)
- KTRs on longterm steroids or at high risk for osteoporosis should undergo DEXA scanning if eGFR >30 ml/min/1.73m² (2D)
- treatment should be according the RCP guidelines for steroid-induced osteoporosis (2D)

Audit Measures

- Prevalence of KTRs on corticosteroids
- Frequency of bisphosphonate usage amongst KTRs
- Incidence of fractures amongst KTRs

Rationale

All KTRs have a complex bone disorder whereby the effects of immunosuppression are superimposed on an underlying Chronic Kidney Disease Mineral and Bone Disorder (CKD-MBD). Any guidance should be used in conjunction with existing guidelines for CKD-MBD⁵¹. The risk of fractures after renal transplantation is high but there is no accurate way to predict fracture risk. Clinical tools have not been validated in KTRs. Bone Mineral Density may not reflect the future risk of fracture in KTRs, particularly in those with eGFR <30 mL/min/1.73m²²⁰³. In addition bisphosphonates are contraindicated in subjects with eGFR <30 mL/min/1.73m². There is evidence that treatment with calcium and vitamin D derivatives attenuates post-transplant bone loss and maintains bone mineral density without excess hypercalcaemia²⁰³⁻²⁰⁵. Corticosteroids seem to be the principal determinant of bone turnover and bone volume so it seems logical to target interventions towards reduction or withdrawal of these drugs²⁰⁵. There are numerous guidelines for corticosteroid induced osteoporosis including those of the Royal College of Physicians and it seems reasonable to follow them^{205,206}. Newer agents such as denosumab, a monoclonal antibody, which inhibits RANK ligand, seem to be safe in renal impairment with the caveat that serum calcium needs to be closely monitored, as the risk of hypocalcaemia is increased²⁰⁷. One randomised control trial of denosumab in the KTR showed some benefit on preservation of bone mineral density, but at the expense of greater numbers of urinary tract infection.²⁰⁸

References

203. Josephson MA, Schumm LP, Chiu MY, Marshall C, Thistlethwaite JR, Sprague SM. Calcium and calcitriol prophylaxis attenuates posttransplant bone loss. *Transplantation* 2004;78:1233-6.
204. Torres A, Garcia S, Gomez A, et al. Treatment with intermittent calcitriol and calcium reduces bone loss after renal transplantation. *Kidney Int* 2004;65:705-12.
- 205 Rizzoli R, Biver E. Glucocorticoid-induced osteoporosis: who to treat with what agent? *Nature reviews Rheumatology* 2015;11:98-109.
206. Compston J. Management of glucocorticoid-induced osteoporosis. *Nature reviews Rheumatology* 2010;6:82-8.
207. Block GA, Bone HG, Fang L, Lee E, Padhi D. A single-dose study of denosumab in patients with various degrees of renal impairment. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* 2012;27:1471-9.
208. Bonani M, Frey D, Brockmann J, Fehr T, Mueller TF, Saleh L, von Eckardstein A, Graf N, Wüthrich RP. Effect of Twice-Yearly Denosumab on Prevention of Bone Mineral Density Loss in De Novo Kidney Transplant Recipients: A Randomized Controlled Trial. *Am J Transplant.* 2016 Jun;16(6):1882-91

Guideline 9.2 KTR: Tertiary hyperparathyroidism

We suggest:

- Severe hyperparathyroidism should be treated prior to transplantation (2D)
- Cinacalcet can be used in KTRs (2C)
- We suggest treatment should be the same as for other patients with CKD (2D)

Audit Measures

- Incidence of hyperparathyroidism
- Incidence of parathyroidectomy
- Usage of cinacalcet

Rationale

Post-transplant hyperparathyroidism is a complex entity that may represent a true high bone turnover state but also low bone turnover²⁰⁹. Dietary intake of phosphate, longstanding phosphate retention whilst on dialysis treatment, followed by sometimes profound urinary phosphate losses in the early post transplant period all contribute to the complexity of managing this condition. In the latter case the suppression of parathyroid hormone (PTH) secretion may lead to adynamic bone disease and the only certain way to distinguish between the two types of mineral and bone disorder (MBD-CKD) is by bone biopsy. There are contradictory data on the effect of parathyroidectomy post transplantation but it seems sensible to treat severe hyperparathyroidism prior to transplantation^{210,211}. In KTRs, cinacalcet may be used; it successfully corrects hypercalcaemia and elevated PTH level, seems to improve bone mineral density, and may also have an antihypertensive effect²¹²⁻²¹⁴. Caution should be exercised with high doses²¹⁵. In hyperparathyroidism refractory to pharmacological agents, parathyroidectomy also provides a benefit but should only be pursued when potential risks of such a procedure are considered.

References

209. Borchhardt K, Sulzbacher I, Benesch T, Fodinger M, Sunder-Plassmann G, Haas M. Low-turnover bone disease in hypercalcemic hyperparathyroidism after kidney transplantation. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2515-21.
210. Evenepoel P, Claes K, Kuypers D, Maes B, Vanrenterghem Y. Impact of parathyroidectomy on renal graft function, blood pressure and serum lipids in kidney transplant recipients: a single centre study. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2005;20:1714-20.
211. Kandil E, Florman S, Alabbas H, et al. Exploring the effect of parathyroidectomy for tertiary hyperparathyroidism after kidney transplantation. *The American journal of the medical sciences* 2010;339:420-4.
212. Bergua C, Torregrosa JV, Fuster D, Gutierrez-Dalmau A, Oppenheimer F, Campistol JM. Effect of cinacalcet on hypercalcemia and bone mineral density in renal transplanted patients with secondary hyperparathyroidism. *Transplantation* 2008;86:413-7.
213. Srinivas TR, Schold JD, Womer KL, et al. Improvement in hypercalcemia with cinacalcet after kidney transplantation. *Clinical journal of the American Society of Nephrology : CJASN* 2006;1:323-6.
214. Zitt E, Woess E, Mayer G, Lhotta K. Effect of cinacalcet on renal electrolyte handling and systemic arterial blood pressure in kidney transplant patients with persistent hyperparathyroidism. *Transplantation* 2011;92:883-9.
215. Serra AL, Braun SC, Starke A, et al. Pharmacokinetics and pharmacodynamics of cinacalcet in patients with hyperparathyroidism after renal transplantation. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2008;8:803-10.

Guideline 9.3 Gout

Guideline 9.3.1 Treatment of gout

We recommend that neither allopurinol nor febuxostat should not be administered with azathioprine (1B)

Guideline 9.3.2 –KTR: Treatment of gout

We suggest:

- Hyperuricaemia should be treated when associated with gout, tophi or uric acid stones (2D)
- Non steroidal anti-inflammatory drugs (NSAIDs) should be avoided in KTRs (2D)
- Episodes of gout may be treated with brief courses of oral prednisolone (2D)
- Colchicine is an effective treatment for gout in KTRs (2D)

Audit Measure

- Frequency of gout and hyperuricaemia amongst KTRs

Rationale

Gout is common after transplantation and may cause significant morbidity. Hyperuricaemia increases the risk of gout and may also be linked with increased rates of cardiovascular disease²¹⁶. In CKD, febuxostat may have superior urate lowering effects compared with allopurinol²¹⁷. This observation is also borne out in the transplant population where febuxostat appears to effectively lower urate levels without adversely affecting renal transplant function, although long term outcome data are lacking^{218,219}. Important drug interactions alter the strategy for managing gout in KTRs. In particular, use of concomitant use of such agents with azathioprine can precipitate potentially fatal blood dyscrasias. CNIs are associated with higher uric acid levels and may contribute to the development of gout.

References

216. Abbott KC, Kimmel PL, Dharnidharka V, Oglesby RJ, Agodoa LY, Caillard S. New-onset gout after kidney transplantation: incidence, risk factors and implications. *Transplantation* 2005;80:1383-91.
217. Sezai A, Soma M, Nakata KI, et al. Comparison of febuxostat and allopurinol for hyperuricemia in cardiac surgery patients with chronic kidney disease (NU-FLASH trial for CKD). *Journal of cardiology* 2015;66(4):289-303.
218. Sofue T, Inui M, Hara T, et al. Efficacy and safety of febuxostat in the treatment of hyperuricemia in stable kidney transplant recipients. *Drug design, development and therapy* 2014;8:245-53.
219. Tojimbara T, Nakajima I, Yashima J, Fuchinoue S, Teraoka S. Efficacy and safety of febuxostat, a novel nonpurine selective inhibitor of xanthine oxidase for the treatment of hyperuricemia in kidney transplant recipients. *Transplantation proceedings* 2014;46:511-3.

Guideline 9.4 –KTR: Calcineurin inhibitor bone pain

We suggest:

- Reducing or withdrawing CNIs should be considered in KTRs with intractable bone pain (2D)
- Dihydropyridine calcium antagonists also may be beneficial (2D)

Rationale

It has become increasingly recognised that CNIs may cause bone pain, which preferentially affects bones in the lower legs^{220,221}. Bone marrow oedema can be demonstrated on magnetic resonance imaging (MRI) scanning and treatment involves reducing CNI levels and the use of dihydropyridine calcium antagonists²²².

References

220. Collini A, De Bartolomeis C, Barni R, Ruggieri G, Bernini M, Carmellini M. Calcineurin-inhibitor induced pain syndrome after organ transplantation. *Kidney Int* 2006;70:1367-70.

221. Grotz WH, Breitenfeldt MK, Braune SW, et al. Calcineurin-inhibitor induced pain syndrome (CIPS): a severe disabling complication after organ transplantation. *Transplant international : official journal of the European Society for Organ Transplantation* 2001;14:16-23.

222. Elder GJ. From marrow oedema to osteonecrosis: common paths in the development of post-transplant bone pain. *Nephrology* 2006;11:560-7.

10. Kidney Transplant Recipient (KTR): Haematological Complications (Guidelines 10.1–10.3)

Guideline 10.1 –KTR: Anaemia

We suggest that chronic anaemia should be managed in the same way as other patient with CKD (2D)

Audit Measure

- Prevalence of anaemia amongst KTRs

Rationale

Anaemia is common in the KTR population, with reports suggesting a prevalence of 20-30% of KTRs, and may be associated with poor outcomes^{111,22}. Its aetiology is multifactorial and may include relative erythropoietin deficiency related to suboptimal graft function, haematinic deficiency, post-transplant infection, and adjunctive antimicrobial therapy (valganciclovir, co-trimoxazole). It may be exacerbated by immunosuppressant therapy, especially antiproliferative agents, and these may be tapered or stopped to improve haemoglobin levels. Management should be similar to other patients with CKD²²⁴. There is one small randomized controlled trial suggesting benefit with haemoglobin correction using an erythropoiesis-stimulating agent²²⁵. However, this is in contrast to larger similar studies in the non-transplant CKD patients where haemoglobin correction to similar levels was associated with adverse outcomes such as increased stroke risk²²⁶.

References

224. Malyszko J, Oberbauer R, Watschinger B. Anemia and erythrocytosis in patients after kidney transplantation. *Transplant international : official journal of the European Society for Organ Transplantation* 2012;25:1013-23.

224. Druke TB, Parfrey PS. Summary of the KDIGO guideline on anemia and comment: reading between the (guide)line(s). *Kidney Int* 2012;82:952-60.

225. Choukroun G, Kamar N, Dussol B, et al. Correction of postkidney transplant anemia reduces progression of allograft nephropathy. *Journal of the American Society of Nephrology : JASN* 2012;23:360-8.

226. Pfeffer MA, Burdmann EA, Chen CY, et al. A trial of darbepoetin alfa in type 2 diabetes and chronic kidney disease. *The New England journal of medicine* 2009;361:2019-32.

Guideline 10.2 – KTR: – Polycythaemia

We recommend that initial treatment should be with angiotensin converting enzyme inhibitors (ACEIs) or with angiotensin receptor blockers (ARBs) (1C)

Guideline 10.3 – KTR: – Polycythaemia

We suggest:

- Haemoglobin levels should be monitored at every clinic visit (2D)
- Treatment should be initiated if the haematocrit or packed cell volume exceeds 52% in men and 49% in women (2D)
- Venesection may be used in refractory cases (2D)

Audit Measure

- Prevalence of post transplant erythrocytosis amongst KTRs

Rationale

Post-transplant polycythaemia (erythrocytosis) is common after renal transplantation and may be associated with significant morbidity and mortality^{227,228}. The overall effect on long term outcome is less clear, although it is likely that it increases risk of thrombotic events²²⁹. Studies have shown that ACEIs and ARBs are associated with a drop in haematocrit in KTRs²²¹.

References

227. Vlahakos DV, Marathias KP, Agroyannis B, Madias NE. Posttransplant erythrocytosis. *Kidney Int* 2003;63:1187-94.
228. Kiberd BA. Post-transplant erythrocytosis: a disappearing phenomenon? *Clinical transplantation* 2009;23:800-6.
229. Yildiz A, Cine N, Akkaya V, et al. Comparison of the effects of enalapril and losartan on posttransplantation erythrocytosis in renal transplant recipients: prospective randomized study. *Transplantation* 2001;72:542-4.

11. Kidney Transplant Recipient (KTR): Reproductive issues (Guidelines 11.1-11.5)

Guideline 11.1 – KTR: Conception and contraception (female).

We recommend that MPA-containing immunosuppressant drugs should be stopped prior to conception and replaced appropriately (1A)

Guideline 11.2 – KTR: Conception and contraception (female)

We suggest:

- KTRs should wait for one year after transplant and have stable function before attempting conception (2C)
- Counselling regarding fertility and reproduction should be offered to female KTRs and their partners either prior to transplantation or soon afterwards (2D)
- m-TORi should be stopped prior to conception and replaced as appropriate (2D)
- Pregnancy should be managed jointly with an Obstetrics department with experience of care of KTRs (2D)
- We suggest that KTRs receive aspirin 75mg daily to reduce the risk of pre-eclampsia from 12 weeks gestation until birth of the baby unless there are contraindications (2C)
- The risks and benefits of breastfeeding should be discussed (2D)
- Contraception advice should be similar to the general population (2D)

Guideline 11.3 – KTR: Conception and contraception (male)

We recommend:

- Male KTRs are advised that MPA containing compounds have theoretical teratogenic potential in men taking these agents (1D)
- KTRs should be advised that m-TORi reduce the male sperm count and counselled accordingly (1C)

Guideline 11.4 – KTR: Conception and contraception (male)

We suggest:

- All immunosuppressive drugs other than m-TORi can be used in male KTRs. Advice for MPA is as Guideline 11.3 (2D)
- The decision to continue MPA containing compounds in a male KTR wishing to conceive should balance a the risk of theoretical teratogenicity against the risk of rejection on changing from MPA to azathioprine (2D)
- Men on m-TORi who wish to conceive should discontinue these agents prior to conception and replace them as appropriate (2D)
- Men who wish to maintain fertility should avoid m-TORi or bank sperm prior to starting these drugs (2D)

Audit Measure

- Pregnancy rates and outcomes should be monitored

Rationale

Female fertility returns rapidly after successful renal transplantation and KTRs and their partners need to be counselled about potential pregnancy²³⁰. Pregnancies in KTRs should be deemed above average risk with increased rates of maternal hypertension, pre-eclampsia, prematurity, low birth weight and caesarean section^{231,232}. The risk of pregnancy to allograft function is probably small, particularly with good baseline function²³¹. Immunosuppressive drugs can all have effects on the foetus and appropriate caution should be exercised. Sirolimus and MPA compounds are teratogenic and should be avoided in females^{232,233}. Recent advice from the MHRA also advises that MPA compounds should be avoided in male KTRs prior to conception as there is a theoretical risk of teratogenicity associated with males fathering children whilst receiving these drugs²³⁴. This advice does not appear to be confirmed in registry studies detailing outcomes in pregnancies fathered by KTRs²³⁴. Therefore, in men, any theoretical risk of foetal abnormality associated with use of MPA compounds, should be balanced against the risk of rejection or decline in graft function associated with changing immunosuppression. Further information on this issue has been generated by the Renal Association's Pregnancy and Chronic Kidney Disease Rare Disease Group and is available on the website RareRenal.org: http://bit.ly/Mycophenolate_rec

Alternative immunosuppression should therefore be considered prior to conception. All immunosuppressants are excreted in breast milk, albeit in tiny quantities, and are usually contraindicated. However, toxicity has not been reported after breastfeeding with ciclosporin, prednisolone, azathioprine and tacrolimus^{228,229}. Aspirin is now widely recommended for the prevention of pre-eclampsia in high-risk populations, and whilst there are no specific data to support its use in KTRs, it seems reasonable to follow this approach, adopted from other guidelines^{235,236}. Outcomes of pregnancies fathered by male KTRs are similar to the general population although reporting bias is difficult to account for^{237, 238}.

There are very little data surrounding the use of female contraception in KTRs and so it seems sensible to extrapolate from the general population with similar cautions and contraindications²³⁹. A number of hypothetical risks associated with specific forms of contraception have not been confirmed in observational studies, though the data quality is poor²⁴⁰.

Whilst ESRD is a state of reduced fertility, the rates of female subfertility and infertility in KTRs are largely unknown. There are a small number of case reports of successful outcomes following in vitro fertilisation in KTRs, albeit with similar complications as seen in KTRs with conventional pregnancy²⁴¹. Further research is needed in this area. Sirolimus and presumably other m-TORi are associated with oligospermia, which appears to be reversible on cessation of treatment²⁴⁰⁻²⁴².

References

230. McKay DB, Josephson MA. Pregnancy in recipients of solid organs--effects on mother and child. *The New England journal of medicine* 2006;354:1281-93.
231. Bramham K, Nelson-Piercy C, Gao H, et al. Pregnancy in renal transplant recipients: a UK national cohort study. *Clinical journal of the American Society of Nephrology : CJASN* 2013;8:290-8.
232. Coscia LA, Constantinescu S, Moritz MJ, et al. Report from the National Transplantation Pregnancy Registry (NTPR): outcomes of pregnancy after transplantation. *Clinical transplants* 2010;65-85.
233. Anderka MT, Lin AE, Abuelo DN, Mitchell AA, Rasmussen SA. Reviewing the evidence for mycophenolate mofetil as a new teratogen: case report and review of the literature. *American journal of medical genetics Part A* 2009;149A:1241-8.
234. Sifontis NM, Coscia LA, Constantinescu S, Lavelanet AF, Moritz MJ, Armenti VT. Pregnancy outcomes in solid organ transplant recipients with exposure to mycophenolate mofetil or sirolimus. *Transplantation* 2006;82:1698-702.
235. Morken NH, Diaz-Garcia C, Reisaeter AV, et al. Obstetric and neonatal outcome of pregnancies fathered by males on immunosuppression after solid organ transplantation. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2015;15:1666-73.
236. Bramham K, Chusney G, Lee J, Lightstone L, Nelson-Piercy C. Breastfeeding and tacrolimus: serial monitoring in breast-fed and bottle-fed infants. *Clinical journal of the American Society of Nephrology : CJASN* 2013;8:563-7.
237. Watnick S, Rueda J. Reproduction and contraception after kidney transplantation. *Current opinion in obstetrics & gynecology* 2008;20:308-12.
238. Duley L, Henderson-Smart DJ, Meher S, King JF. Antiplatelet agents for preventing pre-eclampsia and its complications. *The Cochrane database of systematic reviews* 2007:CD004659.
239. Excellence NfC. Hypertension in pregnancy: The management of hypertensive disorders during pregnancy (clinical guideline 107). 2010.
240. Armenti VT, Daller JA, Constantinescu S, et al. Report from the National Transplantation Pregnancy Registry: outcomes of pregnancy after transplantation. *Clinical transplants* 2006:57-70.
241. Paulen ME, Folger SG, Curtis KM, Jamieson DJ. Contraceptive use among solid organ transplant patients: a systematic review. *Contraception* 2010;82:102-12.
242. Norrman E, Bergh C, Wennerholm UB. Pregnancy outcome and long-term follow-up after in vitro fertilization in women with renal transplantation. *Human reproduction* 2015;30:205-13.
243. Bererhi L, Flamant M, Martinez F, Karras A, Thervet E, Legendre C. Rapamycin-induced oligospermia. *Transplantation* 2003;76:885-6.
244. Deutsch MA, Kaczmarek I, Huber S, et al. Sirolimus-associated infertility: case report and literature review of possible mechanisms. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2007;7:2414-21.
245. Huyghe E, Zairi A, Nohra J, Kamar N, Plante P, Rostaing L. Gonadal impact of target of rapamycin inhibitors (sirolimus and everolimus) in male patients: an overview. *Transplant international : official journal of the European Society for Organ Transplantation* 2007;20:305-11.

Guideline 11.5 – KTR: Sexual function

We suggest:

- Men should be counselled about the possible risks of impotence following transplantation surgery that involves the internal iliac artery (2D)
- Specific enquiry should be made regarding sexual dysfunction, preferably at an annual review clinic (2D)
- Care pathways for dealing with sexual dysfunction should be established (2D)
- Sildenafil is safe and effective in male KTRs not taking nitrates (2D)

Audit Measures

- Prevalence of sexual dysfunction in the transplant clinic

Rationale

Sexual dysfunction is very common in both men and women with advanced CKD and manifests with decreased libido and erectile dysfunction. These problems are often improved after successful renal transplantation but remain common²⁴⁶⁻²⁴⁸. Sildenafil is safe and may be effective for erectile dysfunction in KTRs²⁴⁹.

References

246. Filocamo MT, Zanazzi M, Li Marzi V, et al. Sexual dysfunction in women during dialysis and after renal transplantation. *The journal of sexual medicine* 2009;6:3125-31.
247. Malavaud B, Rostaing L, Rischmann P, Sarramon JP, Durand D. High prevalence of erectile dysfunction after renal transplantation. *Transplantation* 2000;69:2121-4.
248. Raggi MC, Siebert SB, Friess H, Schremmer-Danninger E, Thorban S, Dinkel A. Sexual and relationship functioning before and after renal transplantation: a descriptive study with patients and partners. *Scandinavian journal of urology and nephrology* 2012;46:431-6.
249. Sharma RK, Prasad N, Gupta A, Kapoor R. Treatment of erectile dysfunction with sildenafil citrate in renal allograft recipients: a randomized, double-blind, placebo-controlled, crossover trial. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2006;48:128-33.

5. Lay Summary

These guidelines cover the care of patients from the period following kidney transplantation until the transplant is no longer working or the patient dies. During the early phase prevention of acute rejection and infection are the priority. After around 3-6 months, the priorities change to preservation of transplant function and avoiding the long-term complications of immunosuppressive medication (the medication used to suppress the immune system to prevent rejection). The topics discussed include organization of outpatient follow up, immunosuppressive medication, treatment of acute and chronic rejection, and prevention of complications. The potential complications discussed include heart disease, infection, cancer, bone disease and blood disorders. There is also a section on contraception and reproductive issues.

Immediately after the introduction there is a statement of all the recommendations. These recommendations are written in a language that we think should be understandable by many patients, relatives, carers and other interested people. Consequently we have not reworded or restated them in this lay summary. They are graded 1 or 2 depending on the strength of the recommendation by the authors, and A-D depending on the quality of the evidence that the recommendation is based on.

6. Acknowledgements

This document has been externally reviewed by key stake holders according to the process described in the Clinical Practice Guidelines Development Policy Manual. In addition we are particularly grateful to the following individuals for their review and comments; Peter Andrews, Colin Geddes, Adam MacLean, Will McKane, Michael Robson (Consultant Nephrologists), Lisa Burnapp (Consultant Nurse), Derek Manas, Lorna Marson (Consultant Surgeons), Andrea Devaney (UK Renal Pharmacy Group).

Appendix

Abbreviations used in these guidelines

ACEI	angiotensin converting enzyme inhibitor
ACR	albumin:creatinine ratio
ALERT	Assessment of LEscol in Renal Transplantation
AMR	antibody mediated rejection
AMR	antibody mediated rejection
APKD	adult polycystic kidney disease
ARB	angiotensin receptor blocker
ATG	Anti-thymocyte globulin
ATIIR	angiotensin two receptor
AUC	area under the curve
BCG	Bacillus Calmette–Guérin
BHSI	British Society for Histocompatibility and Immunogenetics
BKN	BK virus nephropathy
BMI	body mass index
BPAR	biopsy proven acute rejection
BTS	British Transplant Society
C ₀	trough concentration
C ₂	concentration at 2 hours post dose
CAD	coronary artery disease
CAMR	chronic antibody mediated rejection
CIT	cold ischaemic time
CKD	chronic kidney disease
CKD-MBD	chronic kidney disease mineral bone disorder
CMV	cytomegalovirus
CNI(s)	calcineurin inhibitor(s)
CRF	calculated reaction frequency
CVD	cardiovascular disease
D-	donor negative
D+	donor positive
DCD	donation after circulatory death
DEXA	Dual-energy X-ray absorptiometry
DM	diabetes mellitus
DNA	Did not attend
DNA	deoxyribonucleic acid
DoH	Department of Health
DSA	donor specific antibody
EBV	Epstein Barr Virus
eGFR	estimated glomerular filtration rate
ESRD	end stage renal disease
GRADE	Grades of Recommendation, Assessment, Development and Evaluation
GT	glucose tolerance
HBsAb	hepatitis B surface antibody
HCV	hepatitis C virus
HDL	high-density lipoprotein
HEV	hepatitis E virus
HIV	human immunodeficiency virus
HSV	herpes simplex virus
HT	hypertension
IL2-RA	interleukin-2 receptor antagonists
JBS	Joint British Societies

KDIGO	Kidney Disease Improving Global Outcomes
KTR(s)	kidney transplant recipient(s)
LDL	low density lipoprotein
m-TORi(s)	mammalian target of rapamycin inhibitor(s)
MI	myocardial infarction
MMF	mycophenolate mofetil
MPA	mycophenolic acid
NDSA	new donor specific antibody
NHSCSP	National Health Service Cancer Screening Programme
NICE	National Institute for Clinical Excellence
NMSC	non-melanoma skin cancer
NSAIDs	non-steroidal anti-inflammatory drugs
PCP	pneumocystis pneumonia
PCR	polymerase chain reaction
PCR	protein:creatinine ratio
PML	Progressive multifocal leukoencephalopathy
PRA	panel reactive antibody
PTDM	post transplant diabetes mellitus
PTH	parathyroid hormone
PTLD	post transplant lymphoproliferative disorder
R-	recipient negative
R+	recipient positive
RANK	Receptor Activator of Nuclear Factor κ B
RAS	renin angiotensin system
RCP	Royal College of Physicians
RCT	randomised controlled trial
RNA	ribonucleic acid
RRT	renal replacement therapy
SCAR	subclinical acute rejection
TDAs	T-cell (lymphocyte) Depleting Antibodies
VHL	Von Hippel–Lindau
VZV	varicella zoster virus